

Rabbit EDA Polyclonal Antibody | anti-EDA antibody

EDA Polyclonal Antibody


CATALOG # AAA10715	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
------------------------------	-----------------------	--	-------------------------------------

PRODUCT SYNONYMS


EDA; Polyclonal Antibody; EDA Polyclonal Antibody; ED1; HED; EDA1; EDA2; ODT1; XHED; XLHED; ED1-A1; ED1-A2; STHAGX1; anti-EDA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	391
Applicable Applications for anti-EDA antibody	WB (Western Blot)
Application Notes	WB: 1:500 - 1:2000
Species	Human
Route	Synthetic Peptide
Immunogen	A synthetic peptide of human EDA
Calculated Molecular Weight	49kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse heart using EDA Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse skeletal muscle using EDA Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse liver using EDA Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded rat pancreas using EDA Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded rat liver using EDA Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human uterine cancer using EDA Antibody at dilution of 1:200 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-EDA antibody	The protein encoded by this gene is a type II membrane protein that can be cleaved by furin to produce a secreted form. The encoded protein, which belongs to the tumor necrosis factor family, acts as a homotrimer and may be involved in cell-cell signaling during the development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia, anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene.
--	---

Product Categories/Family for anti-EDA antibody	Polyclonal
---	------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	6166135
NCBI GeneID	1896
UniProt Accession #	Q92838; O75910; Q5JS00; Q5JUM7; Q9UP77; Q9Y6L0; Q9Y6L1; Q9Y6L2; A0AUZ2; A2A337; B7ZLU2; B7ZLU4
Molecular Weight	391
NCBI Official Full Name	Ectodysplasin-A
NCBI Official Synonym Full Names	ectodysplasin A
NCBI Official Symbol	EDA
NCBI Official Synonym Symbols	ED1; HED; EDA1; EDA2; HED1; ODT1; XHED; ECTD1; XLHED; ED1-A1; ED1-A2; EDA-A1; EDA-A2; STHAGX1
NCBI Protein Information	ectodysplasin-A; oligodontia 1; X-linked anhidrotic ectodermal dysplasia protein
UniProt Protein Name	Ectodysplasin-A
UniProt Gene Name	EDA
UniProt Synonym Gene Names	ED1; EDA2; EDA protein
UniProt Entry Name	EDA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/eda-antibody-western-blot-elisa/10715>