

Mouse anti-Human CK17 Antibody | anti-CK17 antibody

CK17 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA108846	Mouse	Human	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CK17; CK17 Antibody; PC; K17; PC2; PCHC1; KRT17; anti-CK17 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Isotype	IgG2b
Clone Number	3B12
Form/Format	Ascitic fluid containing 0.03% sodium azide.
Sequence Length	432
Applicable Applications for anti-CK17 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Purified recombinant fragment of CK17 expressed in E. Coli.
Preparation and Storage	Store at 4 degree C, for long term storage, store at -20 degree C

IMAGES



IHC (Immunohistochemistry)

Figure 2. Immunohistochemical analysis of paraffin-embedded human lung cancer (A), endometrial carcinoma (B), sublingual gland (C) and esophagus (D) tissues using CK17 mouse mAb with DAB staining.



WB (Western Blot)

Figure 1. Western blot analysis using CK17 mouse mAb against Hela (1), MCF-7 (2) and A431 (3) cell lysate.

ADDITIONAL INFORMATION

Related Product Information for anti-CK17 antibody	CK17, also known as KRT17, it is the type I intermediate filament chain keratin 17. It is found in nail beds, hair follicles, sebaceous glands, and other epidermal appendages. Mutations in this gene lead to Jackson-Lawler type pachyonychia congenita and steatocystoma multiplex. May play a role in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial "stem cells". May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation. Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state. Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway. Involved in tissue repair.
Product Categories/Family for anti-CK17 antibody	Autophagy antibody; Cancer; Cardiovascular; Cell Biology; Epigenetics & Nuclear Signaling; Developmental Biologys; Immunology; Drug Discovery Products; Metabolism; Neuroscience; Signal Transduction; Stem Cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557701
NCBI GeneID	3872
NCBI Accession #	NP_000413.1
NCBI GenBank Nucleotide #	NM_000422.2
UniProt Accession #	Q04695; Q6IP98; Q8N1P6; A5Z1M9; A5Z1N0; A5Z1N1; A5Z1N2; A6NDV6; A6NKQ2
Molecular Weight	49kDa
NCBI Official Full Name	keratin, type I cytoskeletal 17
NCBI Official Synonym Full Names	keratin 17
NCBI Official Symbol	KRT17
NCBI Official Synonym Symbols	PC; K17; PC2; 39.1; CK-17; PCHC1
NCBI Protein Information	keratin, type I cytoskeletal 17
UniProt Protein Name	Keratin, type I cytoskeletal 17
UniProt Gene Name	KRT17
UniProt Synonym Gene Names	CK-17; K17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-human-antibody/ck17/108846>