

5'-nucleotidase (NT5E) Recombinant Protein | NT5E recombinant protein

Recombinant Human 5'-nucleotidase (NT5E), partial

CATALOG # AAA113569	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
-------------------------------	---


PRODUCT SYNONYMS

5'-nucleotidase (NT5E); Recombinant Human 5'-nucleotidase (NT5E); partial; Ecto-5'-nucleotidase; CD73; NT5E recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	28-547aa; Partial
Sequence	ELTILHTNDV HSRLEQTSSE SSKCVNASRC MGGVARLFTK VQQIRRAEPN VLLLDAGDQY QGTIWFTVYK GAEVAHFMNA LRYDAMALGN HEFDNGVEGL IEPLLEAKF PILSANIKAK GPLASQISGL YLPYKVLPGV DEVVGIVGYT SKETPFLSNP GTNLVFEDEI TALQPEVDKL KTLNVNKIIA LGHSGFEMDK LIAQKVRGVD VVVGHSNTF LYTGNPPSKE VPAGKYPFIV TSDDGRKVPV VQAYAFGKYL GYLKIEFDER GNVISSHGNP ILLNSSIPED PSIKADINKW RIKLDNYSTQ ELGKTIVYLD GSSQSCRFRE CNMGNLICDA MINNNLRHTD EMFWNHVSMC ILNGGGIRSP IDERNNGTIT WENLAAVLPF GGTFDLVQLK GSTLKKA FEH SVHRYGQSTG EFLQVGGIHV VYDLSRKP GD RVVKLDVLC T KCRVPSYDPL KMDEVYKVIL PNFLANGGDG FQMIKDELLR HDSGDQDINV VSTYISKMKV IYPAVEGRIK
Preparation and Storage	Store at -20 degrees C. For long-term storage, store at -20 degrees C or -80 degrees C. Store working aliquots at 4 degrees C for up to one week. Repeated freezing and thawing is not recommended.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for NT5E recombinant protein	Hydrolyzes extracellular nucleotides into membrane permeable nucleosides. Exhibits AMP-, NAD-, and NMN-nucleosidase activities.
Product Categories/Family for NT5E recombinant protein	Epigenetics and Nuclear Signaling

References

Primary structure of human placental 5'-nucleotidase and identification of the glycolipid anchor in the mature form.Misumi Y., Ogata S., Ohkubo K., Hirose S., Ikehara Y.Eur. J. Biochem. 191:563-569(1990) Signal sequence and keyword trap in silico for selection of full-length human cDNAs encoding secretion or membrane proteins from oligo-capped cDNA libraries.Otsuki T., Ota T., Nishikawa T., Hayashi K., Suzuki Y., Yamamoto J., Wakamatsu A., Kimura K., Sakamoto K., Hatano N., Kawai Y., Ishii S., Saito K., Kojima S., Sugiyama T., Ono T., Okano K., Yoshikawa Y., Aotsuka S., Sasaki N., Hattori A., Okumura K., Nagai K., Sugano S., Isogai T.DNA Res. 12:117-126(2005) The DNA sequence and analysis of human chromosome 6.Mungall A.J., Palmer S.A., Sims S.K., Edwards C.A., Ashurst J.L., Wilming L., Jones M.C., Horton R., Hunt S.E., Scott C.E., Gilbert J.G.R., Clamp M.E., Bethel G., Milne S., Ainscough R., Almeida J.P., Ambrose K.D., Andrews T.D., Ashwell R.I.S., Babbage A.K., Bagguley C.L., Bailey J., Banerjee R., Barker D.J., Barlow K.F., Bates K., Beare D.M., Beasley H., Beasley O., Bird C.P., Blakey S.E., Bray-Allen S., Brook J., Brown A.J., Brown J.Y., Burford D.C., Burrill W., Burton J., Carder C., Carter N.P., Chapman J.C., Clark S.Y., Clark G., Clee C.M., Clegg S., Cobley V., Collier R.E., Collins J.E., Colman L.K., Corby N.R., Coville G.J., Culley K.M., Dhami P., Davies J., Dunn M., Earthrowl M.E., Ellington A.E., Evans K.A., Faulkner L., Francis M.D., Frankish A., Frankland J., French L., Garner P., Garnett J., Ghorri M.J., Gilby L.M., Gillson C.J., Glithero R.J., Graffham D.V., Grant M., Gribble S., Griffiths C., Griffiths M.N.D., Hall R., Halls K.S., Hammond S., Harley J.L., Hart E.A., Heath P.D., Heathcott R., Holmes S.J., Howden P.J., Howe K.L., Howell G.R., Huckle E., Humphray S.J., Humphries M.D., Hunt A.R., Johnson C.M., Joy A.A., Kay M., Keenan S.J., Kimberley A.M., King A., Laird G.K., Langford C., Lawlor S., Leongamornlert D.A., Leversha M., Lloyd C.R., Lloyd D.M., Loveland J.E., Lovell J., Martin S., Mashreghi-Mohammadi M., Maslen G.L., Matthews L., McCann O.T., McLaren S.J., McLay K., McMurray A., Moore M.J.F., Mullikin J.C., Niblett D., Nickerson T., Novik K.L., Oliver K., Overton-Larty E.K., Parker A., Patel R., Pearce A.V., Peck A.I., Phillimore B.J.C.T., Phillips S., Plumb R.W., Porter K.M., Ramsey Y., Ranby S.A., Rice C.M., Ross M.T., Searle S.M., Sehra H.K., Sheridan E., Skuce C.D., Smith S., Smith M., Spraggon L., Squares S.L., Steward C.A., Sycamore N., Tamlyn-Hall G., Tester J., Theaker A.J., Thomas D.W., Thorpe A., Tracey A., Tromans A., Tubby B., Wall M., Wallis J.M., West A.P., White S.S., Whitehead S.L., Whittaker H., Wild A., Willey D.J., Wilmer T.E., Wood J.M., Wray P.W., Wyatt J.C., Young L., Younger R.M., Bentley D.R., Coulson A., Durbin R.M., Hubbard T., Sulston J.E., Dunham I., Rogers J., Beck S.Nature 425:805-811(2003) Isolation and characterization of the promoter of the human 5'-nucleotidase (CD73) -encoding gene.Hansen K.R., Resta R., Webb C.F., Thompson L.F.Gene 167:307-312(1995) Zanolini L., Rosi F., Pagani R., Marinello E. Characterization of soluble vs membrane-bound human placental 5'-nucleotidase.Klemens M.R., Sherman W.R., Holmberg N.J., Ruedi J.M., Low M.G., Thompson L.F.Biochem. Biophys. Res. Commun. 172:1371-1377(1990) Glycoproteomics analysis of human liver tissue by combination of multiple enzyme digestion and hydrazide chemistry.Chen R., Jiang X., Sun D., Han G., Wang F., Ye M., Wang L., Zou H.J. Proteome Res. 8:651-661(2009) Mass-spectrometric identification and relative quantification of N-linked cell surface glycoproteins.Wollscheid B., Bausch-Fluck D., Henderson C., O'Brien R., Bibel M., Schiess R., Aebersold R., Watts J.D.Nat. Biotechnol. 27:378-386(2009) The high-resolution crystal structure of periplasmic Haemophilus influenzae NAD nucleotidase reveals a novel enzymatic function of human CD73 related to NAD metabolism.Garavaglia S., Bruzzzone S., Cassani C., Canella L., Allegrone G., Sturla L., Mannino E., Millo E., De Flora A., Rizzi M.Biochem. J. 441:131-141(2012) An enzyme assisted RP-RPLC approach for in-depth analysis of human liver phosphoproteome.Bian Y., Song C., Cheng K., Dong M., Wang F., Huang J., Sun D., Wang L., Ye M., Zou H.J. Proteomics 96:253-262(2014) Crystal structure of the human ecto-5'-nucleotidase (CD73) insights into the regulation of purinergic signaling.Knapp K., Zebisch M., Pippel J., El-Tayeb A., Muller C.E., Strater N.Structure 20:2161-2173(2012) NT5E mutations and arterial calcifications.St Hilaire C., Ziegler S.G., Markello T.C., Brusco A., Groden C., Gill F., Carlson-Donohoe H., Lederman R.J., Chen M.Y., Yang D., Siegenthaler M.P., Arduino C., Mancini C., Freudenthal B., Stanescu H.C., Zdebik A.A., Chaganti R.K., Nussbaum R.L., Kleta R., Gahl W.A., Boehm M.N. Engl. J. Med. 364:432-442(2011) NT5E mutations that cause human disease are associated with intracellular mistrafficking of NT5E protein.Fausther M., Lavoie E.G., Goree J.R., Baldini G., Dranoff J.A.PLoS ONE 9:E98568-E98568(2014)

NCBI AND UNIPROT INFORMATION

NCBI GI #	325651886
NCBI GeneID	4907
NCBI Accession #	NP_001191742.1
NCBI GenBank Nucleotide #	NM_001204813.1
UniProt Accession #	P21589; O75520; Q5W116; B3KQI8
Molecular Weight	85kD
NCBI Official Full Name	5'-nucleotidase isoform 2 preproprotein
NCBI Official Synonym Full Names	5'-nucleotidase ecto

NCBI Official Symbol	NT5E
NCBI Official Synonym Symbols	NT; eN; NT5; NTE; eNT; CD73; E5NT; CALJA
NCBI Protein Information	5'-nucleotidase
UniProt Protein Name	5'-nucleotidase
UniProt Gene Name	NT5E
UniProt Synonym Gene Names	NT5; NTE; 5'-NT
UniProt Entry Name	5NTD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/nt5e-recombinant-protein-e-coli/113569>