

TNF receptor-associated factor 5 (TRAF5) Recombinant Protein | TRAF5 recombinant protein

Recombinant Human TNF receptor-associated factor 5 (TRAF5)

CATALOG # AAA114431	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
-------------------------------	---


PRODUCT SYNONYMS

TNF receptor-associated factor 5 (TRAF5); Recombinant Human TNF receptor-associated factor 5 (TRAF5); RING finger protein 84; TRAF5 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-557. Full Length
Sequence	MAYSEEHKGM PCGFIRQNSG NSISLDFEPS IEYQFVERLE ERYKCAFCHS VLHNPHTGC GHRFCQHCL SLRELNTVPI CPVDKEVIKS QEVFKDNCCK REVLNLYVYC SNAPGCNAKV ILGRYQDHLQ QCLFQPVQCS NEKCREPVL R KDLKEHLSAS CQFRKEKCLY CKKD VVVINL QNHEENLCPE YPVFCPNNCA KIILKTEVDE HLAVCPAEQ DCPFKHYGCA VTDKRRNLQQ HEHSALREHM RLVLEKNVQL EEQISDLHKS LEQKESKIQQ LAETIKKLEK EFKQFAQLFG KNGSFLPNIQ VFASHIDKSA WLEAQVHQLL QMVNQQQNKF DLRPLMEAVD TVKQKITLLE NNDQRLAVLE EETNKHDTHI NIHKAQLSKN EERFKLLEGT CYNGKLIWKV TDYKMKKREA VDGHTVSIFS QSFYTSRCGY RLCARAYLNG DGSGRGSHLS LYFVVMRGEF DSSLQWPFRQ RVTLMLLDQS GKKNIMETFK PDPNSSSFKR PDGEMNIASG CPRFVAHSV L ENAKNAYIKD DTLFLKVAVD LTDLEDL
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for TRAF5 recombinant protein	Adapter protein and signal transducer that links mbers of the tumor necrosis factor receptor family to different signaling pathways by association with the receptor cytoplasmic domain and kinases. Mediates activation of NF-kappa-B and probably JNK. Ses to be involved in apoptosis. Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR.
---	--

References

Cloning and characterization of a cDNA encoding the human homolog of tumor necrosis factor receptor-associated factor 5 (TRAF5). Mizushima S., Fujita M., Ishida T., Azuma S., Kato K., Hirai M., Otsuka M., Yamamoto T., Inoue J. *Gene* 207:135-140(1998) Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T., Suzuki Y., Nishikawa T., Otsuki T., Sugiyama T., Irie R., Wakamatsu A., Hayashi K., Sato H., Nagai K., Kimura K., Makita H., Sekine M., Obayashi M., Nishi T., Shibahara T., Tanaka T., Ishii S., Yamamoto J., Saito K., Kawai Y., Isono Y., Nakamura Y., Nagahara K., Murakami K., Yasuda T., Iwayanagi T., Wagatsuma M., Shiratori A., Sudo H., Hosoiri T., Kaku Y., Kodaira H., Kondo H., Sugawara M., Takahashi M., Kanda K., Yokoi T., Furuya T., Kikkawa E., Omura Y., Abe K., Kamihara K., Katsuta N., Sato K., Tanikawa M., Yamazaki M., Ninomiya K., Ishibashi T., Yamashita H., Murakawa K., Fujimori K., Tanai H., Kimata M., Watanabe M., Hiraoka S., Chiba Y., Ishida S., Ono Y., Takiguchi S., Watanabe S., Yosida M., Hotuta T., Kusano J., Kanehori K., Takahashi-Fujii A., Hara H., Tanase T.-O., Nomura Y., Togiya S., Komai F., Hara R., Takeuchi K., Arita M., Imose N., Musashino K., Yuuki H., Oshima A., Sasaki N., Aotsuka S., Yoshikawa Y., Matsunawa H., Ichihara T., Shiohata N., Sano S., Moriya S., Momiyama H., Satoh N., Takami S., Terashima Y., Suzuki O., Nakagawa S., Senoh A., Mizoguchi H., Goto Y., Shimizu F., Wakebe H., Hishigaki H., Watanabe T., Sugiyama A., Takemoto M., Kawakami B., Yamazaki M., Watanabe K., Kumagai A., Itakura S., Fukuzumi Y., Fujimori Y., Komiyama M., Tashiro H., Tanigami A., Fujiwara T., Ono T., Yamada K., Fujii Y., Ozaki K., Hirao M., Ohmori Y., Kawabata A., Hikiji T., Kobatake N., Inagaki H., Ikema Y., Okamoto S., Okitani R., Kawakami T., Noguchi S., Itoh T., Shigeta K., Senba T., Matsumura K., Nakajima Y., Mizuno T., Morinaga M., Sasaki M., Togashi T., Oyama M., Hata H., Watanabe M., Komatsu T., Mizushima-Sugano J., Satoh T., Shirai Y., Takahashi Y., Nakagawa K., Okumura K., Nagase T., Nomura N., Kikuchi H., Masuho Y., Yamashita R., Nakai K., Yada T., Nakamura Y., Ohara O., Isogai T., Sugano S. *Nat. Genet.* 36:40-45(2004) Mural R.J., Istrail S., Sutton G., Florea L., Halpern A.L., Mobarry C.M., Lippert R., Walenz B., Shatkay H., Dew I., Miller J.R., Flanigan M.J., Edwards N.J., Bolanos R., Fasulo D., Halldorsson B.V., Hannenhalli S., Turner R., Yooseph S., Lu F., Nusskern D.R., Shue B.C., Zheng X.H., Zhong F., Delcher A.L., Huson D.H., Kravitz S.A., Mouchard L., Reinert K., Remington K.A., Clark A.G., Waterman M.S., Eichler E.E., Adams M.D., Hunkapiller M.W., Myers E.W., Venter J.C. Human TNF receptor-associated factor 5 (TRAF5) cDNA cloning, expression and assignment of the TRAF5 gene to chromosome 1q32. Nakano H., Shindo M., Yamada K., Yoshida M.C., Santee S.M., Ware C.F., Jenkins N.A., Gilbert D.J., Yagita H., Copeland N.G., Okumura K. *Genomics* 42:26-32(1997) The TNF-receptor-associated factor family scaffold molecules for cytokine receptors, kinases and their regulators. Wajant H., Henkler F., Scheurich P. *Cell. Signal.* 13:389-400(2001) Tumor necrosis factor receptor-associated factors (TRAFs). Bradley J.R., Pober J.S. *Oncogene* 20:6482-6491(2001) Herpesvirus entry mediator, a member of the tumor necrosis factor receptor (TNFR) family, interacts with members of the TNFR-associated factor family and activates the transcription factors NF-kappaB and AP-1. Marsters S.A., Ayres T.M., Skubatch M., Gray C.L., Rothe M., Ashkenazi A.J. *Biol. Chem.* 272:14029-14032(1997) ATAR, a novel tumor necrosis factor receptor family member, signals through TRAF2 and TRAF5. Hsu H., Solovyyev I., Colombero A., Elliott R., Kelley M., Boyle W.J. *J. Biol. Chem.* 272:13471-13474(1997) Tumor necrosis factor (TNF) -mediated kinase cascades bifurcation of nuclear factor-kappaB and c-jun N-terminal kinase (JNK/SAPK) pathways at TNF receptor-associated factor 2. Song H.Y., Regnier C.H., Kirschning C.J., Goeddel D.V., Rothe M. *Proc. Natl. Acad. Sci. U.S.A.* 94:9792-9796(1997) Tumor necrosis factor receptor-associated factor (TRAF) 5 and TRAF2 are involved in CD30-mediated NFkappaB activation. Aizawa S., Nakano H., Ishida T., Horie R., Nagai M., Ito K., Yagita H., Okumura K., Inoue J., Watanabe T. *J. Biol. Chem.* 272:2042-2045(1997) ASK1 is essential for JNK/SAPK activation by TRAF2. Nishitoh H., Saitoh M., Mochida Y., Takeda K., Nakano H., Rothe M., Miyazono K., Ichijo H. *Mol. Cell* 2:389-395(1998) CD40-tumor necrosis factor receptor-associated factor (TRAF) interactions regulation of CD40 signaling through multiple TRAF binding sites and TRAF hetero-oligomerization. Pullen S.S., Miller H.G., Everdeen D.S., Dang T.T., Crute J.J., Kehry M.R. *Biochemistry* 37:11836-11845(1998) Activation of OX40 signal transduction pathways leads to tumor necrosis factor receptor-associated factor (TRAF) 2- and TRAF5-mediated NF-kappaB activation. Kawamata S., Hori T., Imura A., Takaori-Kondo A., Uchiyama T. *J. Biol. Chem.* 273:5808-5814(1998) RIP2 is a novel NF-kappaB-activating and cell death-inducing kinase. McCarthy J.V., Ni J., Dixit V.M. *J. Biol. Chem.* 273:16968-16975(1998) TAJ, a novel member of the tumor necrosis factor receptor family, activates the c-Jun N-terminal kinase pathway and mediates caspase-independent cell death. Eby M.T., Jasmin A., Kumar A., Sharma K., Chaudhary P.M. *J. Biol. Chem.* 275:15336-15342(2000) The TRAF family of signal transducers mediates NF-kappaB activation by the TRANCE receptor. Wong B.R., Josien R., Lee S.Y., Vologodskaya M., Steinman R.M., Choi Y.J. *Biol. Chem.* 273:28355-28359(1998) TTRAP, a novel protein that associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor-associated factors (TRAFs), and that inhibits nuclear factor-kappa B activation. Pype S., Declercq W., Ibrahim A., Michiels C., Van Rietschoten J.G.I., Dewulf N., de Boer M., Vandenabeele P., Huylebroeck D., Remacle J.E.J. *J. Biol. Chem.* 275:18586-18593(2000) TAC1 is a TRAF-interacting receptor for TALL-1, a tumor necrosis factor family member involved in B cell regulation. Xia X.-Z., Treanor J., Senaldi G., Khare S.D., Boone T., Kelley M., Theill L.E., Colombero A., Solovyyev I., Lee F., McCabe S., Elliott R., Miner K., Hawkins N., Guo J., Stolina M., Yu G., Wang J., Delaney J., Meng S.-Y., Boyle W.J., Hsu H.J. *Exp. Med.* 192:137-143(2000) B cell maturation protein is a receptor for the tumor necrosis factor family member TALL-1. Shu H.-B., Johnson H. *Proc. Natl. Acad. Sci. U.S.A.* 97:9156-9161(2000) TRAF family proteins link PKR with NF-kappa B activation. Gil J., Garcia M.A., Gomez-Puertas P., Guerra S., Rullas J., Nakano H., Alcamí J., Esteban M. *Mol. Cell. Biol.* 24:4502-4512(2004) Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. Mayya V., Lundgren D.H., Hwang S.-I., Rezaul K., Wu L., Eng J.K., Rodionov V., Han D.K. *Sci. Signal.* 2:RA46-RA46(2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	77404348
NCBI GeneID	7188
NCBI Accession #	NP_001029082.1
NCBI GenBank Nucleotide #	NM_001033910.2
UniProt Accession #	O00463; Q6FHY1; B4DIS9; B4E0A2
Molecular Weight	80.4 kDa
NCBI Official Full Name	TNF receptor-associated factor 5 isoform b
NCBI Official Synonym Full Names	TNF receptor associated factor 5
NCBI Official Symbol	TRAF5
NCBI Official Synonym Symbols	RNF84; MGC:39780
NCBI Protein Information	TNF receptor-associated factor 5
UniProt Protein Name	TNF receptor-associated factor 5
UniProt Gene Name	TRAF5
UniProt Synonym Gene Names	RNF84
UniProt Entry Name	TRAF5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/traf5-recombinant-protein-e-coli/114431>