

Dickkopf-related protein 2 Recombinant Protein | DKK2 recombinant protein

Recombinant Human Dickkopf-related protein 2

CATALOG # AAA116869	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
-------------------------------	---


PRODUCT SYNONYMS

Dickkopf-related protein 2; Recombinant Human Dickkopf-related protein 2; DKK2 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	34-259. Full Length of Mature Protein
Sequence	KLNSIKSSLG GETPGQAANR SAGMYQGLAF GGSKKGKNLG QAYPCSSDKE CEVGRYCHSP HQGSSACMVC RRKKKRCHRD GMCCPSTRCN NGICIPVTES ILTPHIPALD GTRHRDRNHG HYSNHDLGWQ NLGRPHTKMS HIKGHEGDPC LRSSDCIEGF CCARHFWTKI CKPVLHQGEV CTQRKKKGSH GLEIFQR CDC AKGLSCKVWK DATYSSKARL HVCQKI
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for DKK2 recombinant protein	Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease.
Product Categories/Family for DKK2 recombinant protein	Signal Transduction

References

Functional and structural diversity of the human Dickkopf gene family.Krupnik V.E., Sharp J.D., Jiang C., Robison K., Chickering T.W., Amaravadi L., Brown D.E., Guyot D., Mays G., Leiby K., Chang B., Duong T., Goodearl A.D.J., Gearing D.P., Sokol S.Y., McCarthy S.A.Gene 238:301-313(1999) Tanaka S., Sugimachi K., Sugimachi K.The secreted protein discovery initiative (SPDI) , a large-scale effort to identify novel human secreted and transmembrane proteins a bioinformatics assessment.Clark H.F., Gurney A.L., Abaya E., Baker K., Baldwin D.T., Brush J., Chen J., Chow B., Chui C., Crowley C., Currell B., Deuel B., Dowd P., Eaton D., Foster J.S., Grimaldi C., Gu Q., Hass P.E., Heldens S., Huang A., Kim H.S., Klimowski L., Jin Y., Johnson S., Lee J., Lewis L., Liao D., Mark M.R., Robbie E., Sanchez C., Schoenfeld J., Seshagiri S., Simmons L., Singh J., Smith V., Stinson J., Vagts A., Vandlen R.L., Watanabe C., Wieand D., Woods K., Xie M.-H., Yansura D.G., Yi S., Yu G., Yuan J., Zhang M., Zhang Z., Goddard A.D., Wood W.I., Godowski P.J., Gray A.M.Genome Res. 13:2265-2270(2003) Complete sequencing and characterization of 21,243 full-length human cDNAs.Ota T., Suzuki Y., Nishikawa T., Otsuki T., Sugiyama T., Irie R., Wakamatsu A., Hayashi K., Sato H., Nagai K., Kimura K., Makita H., Sekine M., Obayashi M., Nishi T., Shibahara T., Tanaka T., Ishii S., Yamamoto J., Saito K., Kawai Y., Isono Y., Nakamura Y., Nagahari K., Murakami K., Yasuda T., Iwayanagi T., Wagatsuma M., Shiratori A., Sudo H., Hosoiri T., Kaku Y., Kodaira H., Kondo H., Sugawara M., Takahashi M., Kanda K., Yokoi T., Furuya T., Kikkawa E., Omura Y., Abe K., Kamihara K., Katsuta N., Sato K., Tanikawa M., Yamazaki M., Ninomiya K., Ishibashi T., Yamashita H., Murakawa K., Fujimori K., Tanai H., Kimata M., Watanabe M., Hiraoka S., Chiba Y., Ishida S., Ono Y., Takiguchi S., Watanabe S., Yosida M., Hotuta T., Kusano J., Kanehori K., Takahashi-Fujii A., Hara H., Tanase T.-O., Nomura Y., Togiya S., Komai F., Hara R., Takeuchi K., Arita M., Imose N., Musashino K., Yuuki H., Oshima A., Sasaki N., Aotsuka S., Yoshikawa Y., Matsunawa H., Ichihara T., Shiohata N., Sano S., Moriya S., Momiyama H., Satoh N., Takami S., Terashima Y., Suzuki O., Nakagawa S., Senoh A., Mizoguchi H., Goto Y., Shimizu F., Wakebe H., Hishigaki H., Watanabe T., Sugiyama A., Takemoto M., Kawakami B., Yamazaki M., Watanabe K., Kumagai A., Itakura S., Fukuzumi Y., Fujimori Y., Komiyama M., Tashiro H., Tanigami A., Fujiwara T., Ono T., Yamada K., Fujii Y., Ozaki K., Hirao M., Ohmori Y., Kawabata A., Hikiji T., Kobatake N., Inagaki H., Ikema Y., Okamoto S., Okitani R., Kawakami T., Noguchi S., Itoh T., Shigeta K., Senba T., Matsumura K., Nakajima Y., Mizuno T., Morinaga M., Sasaki M., Togashi T., Oyama M., Hata H., Watanabe M., Komatsu T., Mizushima-Sugano J., Satoh T., Shirai Y., Takahashi Y., Nakagawa K., Okumura K., Nagase T., Nomura N., Kikuchi H., Masuho Y., Yamashita R., Nakai K., Yada T., Nakamura Y., Ohara O., Isogai T., Sugano S.Nat. Genet. 36:40-45(2004)

NCBI AND UNIPROT INFORMATION

NCBI GeneID	27123
UniProt Accession #	Q9UBU2; Q9UIU3; A0AVE9; B2R6S7
Molecular Weight	41 kDa
NCBI Official Synonym Full Names	dickkopf WNT signaling pathway inhibitor 2
NCBI Official Symbol	DKK2
NCBI Official Synonym Symbols	DKK-2
NCBI Protein Information	dickkopf-related protein 2
UniProt Protein Name	Dickkopf-related protein 2
UniProt Gene Name	DKK2
UniProt Synonym Gene Names	Dickkopf-2; Dkk-2; hDkk-2
UniProt Entry Name	DKK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.