

Rabbit anti-Human BCMA Monoclonal Antibody | anti-BCMA antibody

Anti-BCMA antibody (DM6), Rabbit mAb


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA11704	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence

PRODUCT SYNONYMS

BCMA; Monoclonal Antibody; Anti-BCMA antibody (DM6); Rabbit mAb; TNFRSF17; CD269; BCM; anti-BCMA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM6
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid; 50% Glycerol; PBS, pH7.4; 0.1% BSA
Applicable Applications for anti-BCMA antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence)
Application Notes	FC:FACS: 1:100
Immunogen	Recombinant human BCMA (Met1-Ala54) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preservative	0.1% Procline 300
Constituents	50% Glycerol; PBS, pH 7.4; 0.1% BSA
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing) Shipping: Blue Ice




IP (Immunoprecipitation)

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.




FCM (Flow Cytometry)

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.




FCM (Flow Cytometry)

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.




Application Data

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.



Application Data

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.



Application Data

Figure 6. Immunoprecipitation analysis. Cellular overexpression lysates (made from HEK293F cells transfected with DYKDDDDK tagged human BCMA full length gene) were pre-incubated with 6 different rabbit DimAb clones and negative control IgG. The immunocomplexes were further pulled down by protein A beads, fractionated, and blotted with mouse anti-DYKDDDDK monoclonal antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-BCMA antibody	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq, Jul 2008]
Product Categories/Family for anti-BCMA antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GeneID	608
Molecular Weight	20kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/bcma-antibody-clone-dm6-flow-cytometry-immunofluorescence-elisa/11704>