

Rabbit anti-Human Cellular tumor antigen p53 Polyclonal Antibody | anti-TP53 antibody

Rabbit anti-human Cellular tumor antigen p53 polyclonal Antibody, FITC conjugate

CATALOG # AAA119558	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	-----------------------	----------------------------	------------------------------

PRODUCT SYNONYMS

Cellular tumor antigen p53; Polyclonal Antibody; Rabbit anti-human Cellular tumor antigen p53 polyclonal Antibody; FITC conjugate; Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53; TP53; P53; anti-TP53 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G Purified
Form/Format	Liquid
Sequence Length	393
Applicable Applications for anti-TP53 antibody	ELISA
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate	FITC
Immunogen	Recombinant human Cellular tumor antigen p53 protein(2-393AA)
Species	Homo sapiens (Human)
Research Area	Cell Biology
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1314 / sc-1311-R / sc-6243
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

ADDITIONAL INFORMATION

Related Product Information for anti-TP53 antibody	Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkl1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediatedapoptosis.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	120407068
NCBI GeneID	7157
NCBI Accession #	NP_000537.3
NCBI GenBank Nucleotide #	NM_000546.5
UniProt Accession #	P04637; Q15086; Q15087; Q15088; Q16535; Q16807; Q16808; Q16809; Q16810; Q16811; Q16848; Q2XN98
NCBI Official Full Name	cellular tumor antigen p53 isoform a
NCBI Official Synonym Full Names	tumor protein p53
NCBI Official Symbol	TP53
NCBI Official Synonym Symbols	P53; BCC7; LFS1; TRP53
NCBI Protein Information	cellular tumor antigen p53
UniProt Protein Name	Cellular tumor antigen p53
UniProt Gene Name	TP53
UniProt Synonym Gene Names	P53
UniProt Entry Name	P53_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cellular-tumor-antigen-p53-antibody-elisa/119558>