

Rabbit anti-Human NPC2 Polyclonal Antibody | anti-NPC2 antibody

NPC2 Antibody

CATALOG # AAA119657	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

NPC2; Polyclonal Antibody; NPC2 Antibody; Epididymal secretory protein E1; HE1; Human epididymis-specific protein 1; He1; Niemann-Pick disease type C2 protein; anti-NPC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G Purified
Sequence Length	151
Applicable Applications for anti-NPC2 antibody	IHC (Immunohistochemistry), ELISA
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate	Non-conjugated
Immunogen	Recombinant human Epididymal secretory protein E1 protein
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-30349 / sc-21470 / sc-30347 / sc-30346 / sc-33776
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of A549 cells using AAA119657 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney tissue using AAA119657 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-NPC2 antibody	Intracellular cholesterol transporter which acts in concert with NPC1 and plays an important role in the egress of cholesterol from the endosomal/lysosomal compartment. Both NPC1 and NPC2 function as the cellular 'tag team duo' (TTD) to catalyze the mobilization of cholesterol within the multivesicular environment of the late endosome (LE) to effect egress through the limiting bilayer of the LE. NPC2 binds unesterified cholesterol that has been released from LDLs in the lumen of the late endosomes/lysosomes and transfers it to the cholesterol-binding pocket of the N-terminal domain of NPC1. Cholesterol binds to NPC1 with the hydroxyl group buried in the binding pocket and is exported from the limiting membrane of late endosomes/ lysosomes to the ER and plasma membrane by an unknown mechanism. The secreted form of NCP2 regulates biliary cholesterol secretion via stimulation of ABCG5/ABCG8-mediated cholesterol transport.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	5453678
NCBI GeneID	10577
NCBI Accession #	NP_006423.1
NCBI GenBank Nucleotide #	NM_006432.3
UniProt Accession #	P61916; Q15668; Q29413; B4DQV7
Molecular Weight	13,416 Da
NCBI Official Full Name	epididymal secretory protein E1
NCBI Official Synonym Full Names	Niemann-Pick disease, type C2
NCBI Official Symbol	NPC2
NCBI Official Synonym Symbols	HE1; EDDM1
NCBI Protein Information	epididymal secretory protein E1
UniProt Protein Name	Epididymal secretory protein E1
UniProt Gene Name	NPC2
UniProt Synonym Gene Names	HE1; He1
UniProt Entry Name	NPC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/npc2-antibody-immunohistochemistry-elisa/119657>