

Rabbit IL33 Polyclonal Antibody | anti-IL33 antibody

Anti-IL33 Picoband antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124549	Rabbit	Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

IL33; Polyclonal Antibody; Anti-IL33 Picoband antibody; Interleukin-33; IL-33; IL33; anti-IL33 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	264
Applicable Applications for anti-IL33 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived rat IL33 recombinant protein (Position: A102-M264).
Subcellular Localization	Nucleus.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES




IHC (Immunohistochemistry)

Figure 4. IHC analysis of IL33 using anti-IL33 antibody (AAA124549). IL33 was detected in paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL33 Antibody (AAA124549) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of IL33 using anti-IL33 antibody (AAA124549). IL33 was detected in paraffin-embedded section of rat lymphaden tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL33 Antibody (AAA124549) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of IL33 using anti-IL33 antibody (AAA124549). IL33 was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL33 Antibody (AAA124549) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of IL33 using anti-IL33 antibody (AAA124549). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat spleen tissue lysates, Lane 2: rat lung tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL33 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for IL33 at approximately 31-37KD. The expected band size for IL33 is at 31KD.

ADDITIONAL INFORMATION

Related Product Information for anti-IL33 antibody	Description: Interleukin 33 (IL-33) is a protein that in humans is encoded by the IL33 gene. The protein encoded by this gene is a cytokine that binds to the IL1RL1/ST2 receptor. The encoded protein is involved in the maturation of Th2 cells and the activation of mast cells, basophils, eosinophils and natural killer cells. Several transcript variants encoding different isoforms have been found for this gene. Protein Function: Cytokine that binds to and signals through the IL1RL1/ST2 receptor which in turn activates NF-kappa-B and MAPK signaling pathways in target cells. Involved in the maturation of Th2 cells inducing the secretion of T-helper type 2-associated cytokines. Also involved in activation of mast cells, basophils, eosinophils and natural killer cells. Acts as a chemoattractant for Th2 cells, and may function as an "alarmin", that amplifies immune responses during tissue injury (By similarity).
Product Categories/Family for anti-IL33 antibody	Cardiovascular; Angiogenesis; Cytokines; Immunology; Innate Immunity; Interleukins; Cardiovascular; Heart; Hypertrophy

NCBI AND UNIPROT INFORMATION

NCBI GI #	62079057
NCBI GeneID	361749
NCBI Accession #	NP_001014188.1
NCBI GenBank Nucleotide #	NM_001014166.1
UniProt Accession #	Q66H70
Molecular Weight	29,486 Da
NCBI Official Full Name	interleukin-33
NCBI Official Synonym Full Names	interleukin 33
NCBI Official Symbol	IL33
NCBI Official Synonym Symbols	RGD1311155
NCBI Protein Information	interleukin-33

UniProt Protein Name	Interleukin-33
UniProt Gene Name	IL33
UniProt Synonym Gene Names	IL-33

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/il33-antibody-western-blot-immunohistochemistry-elisa/124549>