

Rabbit IL17 Polyclonal Antibody | anti-IL17a antibody

Anti-IL17 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124572	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

IL17; Polyclonal Antibody; Anti-IL17 Picoband Antibody; Interleukin-17A; IL-17; IL-17A; Cytotoxic T-lymphocyte-associated antigen 8; CTLA-8; IL17a; Ctl8; IL17; anti-IL17a antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Sequence Length	158
Applicable Applications for anti-IL17a antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived mouse IL17 recombinant protein (Position: A26-A158).
Subcellular Localization	Secreted.
Tissue Specificity	Restricted to a subset of activated T-cells.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of IL17 using anti-IL17 antibody (AAA124572).IL17 was detected in paraffin-embedded section of rat pancreas tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL17 Antibody (AAA124572) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 2. IHC analysis of IL17 using anti-IL17 antibody (AAA124572).IL17 was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL17 Antibody (AAA124572) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of IL17 using anti-IL17 antibody (AAA124572).IL17 was detected in paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IL17 Antibody (AAA124572) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of IL17 using anti-IL17 antibody (AAA124572).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: human Jurkat cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL17 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for IL17 at approximately 27KD. The expected band size for IL17 is at 17KD.

ADDITIONAL INFORMATION

Related Product Information for anti-IL17a antibody	Description: Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. Protein Function: Ligand for IL17RA (PubMed: 17911633). The heterodimer formed by IL17A and IL17F is a ligand for the heterodimeric complex formed by IL17RA and IL17RC (By similarity). Involved in inducing stromal cells to produce proinflammatory and hematopoietic cytokines (By similarity).
Product Categories/Family for anti-IL17a antibody	Immunology; Innate Immunity; Macrophage/Inflamm.; Immunology; Cytokines; interleukins

NCBI AND UNIPROT INFORMATION

NCBI GI #	6754324
NCBI GeneID	16171
NCBI Accession #	NP_034682.1
NCBI GenBank Nucleotide #	NM_010552.3
UniProt Accession #	Q62386; Q60971
NCBI Official Full Name	interleukin-17A
NCBI Official Synonym Full Names	interleukin 17A
NCBI Official Symbol	IL17a
NCBI Official Synonym Symbols	IL17; Ctla8; IL-17; Ctla-8; IL-17A
NCBI Protein Information	interleukin-17A
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	IL17a
UniProt Synonym Gene Names	Ctla8; IL17; IL-17; IL-17A; CTLA-8

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/il17-antibody-western-blot-immunohistochemistry-elisa/124572>