

Rabbit IGFBP2 Polyclonal Antibody | anti-IGFBP2 antibody

Anti-IGFBP2 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124643	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

IGFBP2; Polyclonal Antibody; Anti-IGFBP2 Picoband Antibody; Insulin-like growth factor-binding protein 2; IBP-2; IGF-binding protein 2; IGFBP-2; BP2; IBP2; anti-IGFBP2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Concentration	Add 0.2ml of distilled water will yield a concentration of 500ug/ml. (varies by lot)
Sequence Length	325
Applicable Applications for anti-IGFBP2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived human IGFBP2 recombinant protein (Position: A36-Q325).
Subcellular Localization	Secreted.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemisry)

Figure 3. IHC analysis of IGFBP2 using anti-IGFBP2 antibody (AAA124643).IGFBP2 was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IGFBP2 Antibody (AAA124643) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohiostchemistry)

Figure 2. IHC analysis of IGFBP2 using anti-IGFBP2 antibody (AAA124643).IGFBP2 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-IGFBP2 Antibody (AAA124643) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of IGFBP2 using anti-IGFBP2 antibody (AAA124643).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: human HepG2 whole cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IGFBP2 antigen affinity purified polyclonal antibody at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for IGFBP2 at approximately 35KD. The expected band size for IGFBP2 is at 35KD.

ADDITIONAL INFORMATION

Related Product Information for anti-IGFBP2 antibody	Description: The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins with conserved cysteine residues, which are clustered in the amino- and carboxy-terminal thirds of the molecule. IGFBPs modulate the biological activities of IGF proteins. Some IGFBPs may also have intrinsic bioactivity that is independent of their ability to bind IGF proteins. Post-translational modifications of IGFBPs, including glycosylation, phosphorylation and proteolysis, have been shown to modify the affinities of the binding proteins to IGF. Human IGFBP-2 cDNA encodes a 328 amino acid (aa) residue precursor protein with a putative 39 aa residue signal peptide that is processed to generate the 289 aa residue mature protein. IGFBP-2 contains an integrin receptor recognition sequence (RGD sequence) but lacks potential N-linked glycosylation sites. During development, IGFBP-2 is expressed in a number of tissues. The highest expression level is found in the central nervous system. In adults, high expression levels are also detected in the central nervous system and in a number of reproductive tissues. IGFBP-2 binds preferentially to IGF II, exhibiting a 2-10 fold higher affinity for IGF II than for IGF I. Protein Function: Inhibits IGF-mediated growth and developmental rates. IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.
Product Categories/Family for anti-IGFBP2 antibody	Signal Transduction; Growth Factors/Hormones; insulin/insulin-like; Cancer; Growth Factors; insulin and insulin-like; Invasion/Microenvironment; ECM; extracellular matrix

NCBI AND UNIPROT INFORMATION

NCBI GI #	55925576
NCBI GeneID	3485
NCBI Accession #	NP_000588.2
NCBI GenBank Nucleotide #	NM_000597.2
UniProt Accession #	P18065; Q14619; Q9UCL3
Molecular Weight	34,814 Da
NCBI Official Full Name	insulin-like growth factor-binding protein 2 isoform a
NCBI Official Synonym Full Names	insulin like growth factor binding protein 2
NCBI Official Symbol	IGFBP2
NCBI Official Synonym Symbols	IBP2; IGF-BP53
NCBI Protein Information	insulin-like growth factor-binding protein 2
UniProt Protein Name	Insulin-like growth factor-binding protein 2
UniProt Gene Name	IGFBP2
UniProt Synonym Gene Names	BP2; IBP2; IBP-2; IGF-binding protein 2; IGFBP-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/igfbp2-antibody-western-blot-immunohistochemistry-elisa/124643>