

Mouse TFF3 ELISA Kit | TFF3 elisa kit

Mouse TFF3 PicoKine ELISA Kit

CATALOG #
AAA124993

REACTIVITY
Mouse


PRODUCT SYNONYMS

TFF3; Mouse TFF3 PicoKine ELISA Kit; Mouse TFF3; TFF3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Natural and recombinant Mouse STC2
Sequence Length	94
Samples	Cell Culture Supernates, Serum And Plasma(Heparin)
Detection Range	156pg/ml-10000pg/ml
Sensitivity	<50pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay): Three samples of known concentration were tested on one plate to assess intra-assay precision
Inter-assay Precision	Inter-Assay Precision (Precision accross assays):Three samples of known concentration were tested in separate assays to assess inter-assay precision.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TFF3 elisa kit	Principle of the Assay: The Picokine Mouse STC2 Pre-Coated ELISA (Enzyme-Linked Immunosorbent Assay) kit is a solid phase immunoassay specially designed to measure Mouse STC2 with a 96-well strip plate that is pre-coated with antibody specific for STC2. The detection antibody is a biotinylated antibody specific for STC2. The capture antibody is A monoclonal antibody from rat, the detection antibody is a biotinylated detection polyclonal antibody from goat. The kit contains recombinant Mouse STC2 with immunogen: Expression system for standard: NSO, Immunogen sequence: T25-R296. The kit is analytically validated with ready to use reagents. To measure Mouse STC2, add standards and samples to the wells, then add the biotinylated detection antibody. Wash the wells with PBS or TBS buffer, and add Avidin-Biotin-Peroxidase Complex (ABC-HRP). Wash away the unbounded ABC-HRP with PBS or TBS buffer and add TMB. TMB is substrate to HRP and will be catalyzed to produce a blue color product, which changes into yellow after adding acidic stop solution. The density of the yellow product is linearly propotional to Mouse STC2 in the sample. Read the density of the yellow product in each well using a plate reader, and benchmark the sample wells' readings against the standard curve to determine the concentration of Mouse STC2 in the sample!!Background/Introduction: Stanniocalcin-2 is a protein that in humans is encoded by the STC2 gene. It is mapped to 11; 11 A4. This gene encodes a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The encoded protein has 10 of its 15 cysteine residues conserved among stanniocalcin family members and is phosphorylated by casein kinase 2 exclusively on its serine residues. Its Cterminus contains a cluster of histidine residues which may interact with metal ions. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Constitutive overexpression of human stanniocalcin 2 in mice resulted in pre-and postnatal growth restriction, reduced bone and skeletal muscle growth, and organomegaly. Expression of this gene is induced by estrogen and altered in some breast cancers.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	281485608
NCBI GeneID	7033
NCBI Accession #	NP_003217.3
NCBI GenBank Nucleotide #	NM_003226.3
NCBI Official Full Name	trefoil factor 3
NCBI Official Synonym Full Names	trefoil factor 3
NCBI Official Symbol	TFF3
NCBI Official Synonym Symbols	ITF; P1B; TFI
NCBI Protein Information	trefoil factor 3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.