

Rabbit anti-Human Caspase-8/CASP8 Polyclonal Antibody | anti-CASP8 antibody

Anti-Caspase-8/CASP8 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125947	Rabbit	Human	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

Caspase-8/CASP8; Polyclonal Antibody; Anti-Caspase-8/CASP8 Antibody Picoband; anti-CASP8 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CASP8 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Caspase-8/CASP8 recombinant protein (Position: S217-D479).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of K562 cells using anti-Caspase-8/CASP8 antibody (AAA125947). Overlay histogram showing K562 cells stained with AAA125947 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Caspase-8/CASP8 Antibody (AAA125947, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of Caspase-8/CASP8 using anti-Caspase-8/CASP8 antibody (AAA125947). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: human Raji whole cell lysates, Lane 4: human K562 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Caspase-8/CASP8 antigen affinity purified polyclonal antibody (#AAA125947) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Caspase-8/CASP8 at approximately 55 kDa. The expected band size for Caspase-8/CASP8 is at 55 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CASP8 antibody	CASP8 is also known as CAP4, MACH or MCH5. This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. In addition, this protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined.
Product Categories/Family for anti-CASP8 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	122056474
NCBI GeneID	841
NCBI Accession #	NP_001073593.1
NCBI GenBank Nucleotide #	NM_001080124.1
UniProt Accession #	Q14790; O14676; Q14791; Q14792; Q14793; Q14794; Q14795; Q14796; Q15780; Q15806; Q53TT5; Q8TDI1
Molecular Weight	55,391 Da
NCBI Official Full Name	caspase-8 isoform C
NCBI Official Synonym Full Names	caspase 8, apoptosis-related cysteine peptidase

NCBI Official Symbol	CASP8
NCBI Official Synonym Symbols	CAP4; MACH; MCH5; FLICE; ALPS2B; Casp-8
NCBI Protein Information	caspase-8; FADD-like ICE; MACH-alpha-1/2/3 protein; apoptotic protease Mch-5; MACH-beta-1/2/3/4 protein; apoptotic cysteine protease; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; FADD-homologous ICE/CED-3-like protease; caspase 8, apopto
UniProt Protein Name	Caspase-8
UniProt Gene Name	CASP8
UniProt Synonym Gene Names	MCH5; CASP-8; FLICE; MACH
UniProt Entry Name	CASP8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/caspase-8-casp8-antibody-western-blot-flow-cytometry-elisa/125947>