

Rabbit anti-Human MUC1 Polyclonal Antibody | anti-MUC1 antibody

Anti-MUC1 Antibody Picoband



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125974	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Immunohistochemistry

PRODUCT SYNONYMS

MUC1; Polyclonal Antibody; Anti-MUC1 Antibody Picoband; anti-MUC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MUC1 antibody	ELISA, FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MUC1 recombinant protein (Position: S1098-L1255).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of A431 cells using anti-MUC1 antibody (AAA125974).Overlay histogram showing A431 cells stained with AAA125974 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MUC1 Antibody (AAA125974, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

IHC (Immunohistochemistry)

Figure 1. IHC analysis of MUC1 using anti-MUC1 antibody (AAA125974).MUC1 was detected in a paraffin-embedded section of human rectal moderately differentiated adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MUC1 Antibody (AAA125974) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MUC1 antibody	Mucin 1, cell surface associated (MUC1) or polymorphic epithelial mucin (PEM) is a mucin encoded by the MUC1 gene in humans. This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. It is mapped to 1q22. Mucin 1 is a transmembrane mucin normally expressed on the apical borders of secretory epithelial cells. Overexpression of Mucin 1 is often associated with colon, breast, ovarian, lung and pancreatic cancers. The protein serves a protective function by binding to pathogens and also functions in a cell signaling capacity. Mucin 1 stimulated ESR1-mediated transcription and contributed to estradiol-mediated growth and survival of breast cancer cells. This gene also can suppress pulmonary innate immunity, and its antiinflammatory activity may play an important modulatory role during microbial infection.
Product Categories/Family for anti-MUC1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	67189007
NCBI GeneID	4582
NCBI Accession #	NP_001018016.1
NCBI GenBank Nucleotide #	NM_001018016.2
UniProt Accession #	P15941; P13931; P15942; P17626; Q0VAP6; Q14128; Q14876; Q16437; Q16442; Q16615; B1AVQ8; B1AVR0
Molecular Weight	122,102 Da
NCBI Official Full Name	mucin-1 isoform 2
NCBI Official Synonym Full Names	mucin 1, cell surface associated
NCBI Official Symbol	MUC1
NCBI Official Synonym Symbols	EMA; PEM; PUM; KL-6; MAM6; PEMT; CD227; H23AG; MUC-1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC

NCBI Protein Information	mucin-1; episialin; DF3 antigen; H23 antigen; cancer antigen 15-3; krebs von den Lungen-6; mucin 1, transmembrane; tumor-associated mucin; carcinoma-associated mucin; polymorphic epithelial mucin; peanut-reactive urinary mucin; tumor associated epithelial
UniProt Protein Name	Mucin-1
UniProt Gene Name	MUC1
UniProt Synonym Gene Names	PUM; MUC-1; CA 15-3; KL-6; PUM; PEM; EMA; MUC1-NT; MUC1-alpha; MUC1-beta
UniProt Entry Name	MUC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/muc1-antibody-flow-cytometry-immunohistochemistry-elisa/125974>