

Rabbit E2F1 Polyclonal Antibody | anti-E2F1 antibody

Anti-E2F1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125996	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS

E2F1; Polyclonal Antibody; Anti-E2F1 Antibody Picoband; anti-E2F1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-E2F1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human E2F1 recombinant protein (Position: K89-A328).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of PC-3 cells using anti-E2F1 antibody (AAA125996).Overlay histogram showing PC-3 cells stained with AAA125996 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-E2F1 Antibody (AAA125996, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of E2F1 using anti-E2F1 antibody (AAA125996).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human A431 whole cell lysates,Lane 3: human Caco-2 whole cell lysates,Lane 4: rat brain tissue lysates,Lane 5: rat testis tissue lysates,Lane 6: mouse brain tissue lysates,Lane 7: mouse testis tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-E2F1 antigen affinity purified polyclonal antibody (#AAA125996) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for E2F1 at approximately 70 kDa. The expected band size for E2F1 is at 47 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-E2F1 antibody	Transcription factor E2F1 is a protein that in humans is encoded by the E2F1 gene. The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family of transcription factors appears to play a critical role in the transcription of certain genes required for cell cycle progression. E2F1, the first cloned member of this family, is regulated during the cell cycle at the mRNA level by changes in transcription of the E2F1 gene and at the protein level by complex formation with proteins such as the retinoblastoma gene product(pRB), cyclin A and DP1. E2F1 can override a pRB-induced G1/S block and can behave as an oncogene in certain cells. E2F1 was cloned and was found to contain seven exons. Fluorescence in situ hybridization localized E2F1 to chromosome 20q11. The E2F1 transcription factor can promote proliferation or apoptosis when activated, and is a key downstream target of the retinoblastoma tumour suppressor protein(pRB).
Product Categories/Family for anti-E2F1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	12669911
NCBI GeneID	1869
NCBI Accession #	NP_005216.1
NCBI GenBank Nucleotide #	NM_005225.2
UniProt Accession #	Q01094; Q13143; Q92768
Molecular Weight	46,920 Da
NCBI Official Full Name	transcription factor E2F1
NCBI Official Synonym Full Names	E2F transcription factor 1
NCBI Official Symbol	E2F1

NCBI Official Synonym Symbols	RBP3; E2F-1; RBAP1; RBBP3
NCBI Protein Information	transcription factor E2F1; PBR3; RBAP-1; RBBP-3; PRB-binding protein E2F-1; retinoblastoma-binding protein 3; retinoblastoma-associated protein 1
UniProt Protein Name	Transcription factor E2F1
UniProt Gene Name	E2F1
UniProt Synonym Gene Names	RBBP3; E2F-1; RBAP-1; RBBP-3
UniProt Entry Name	E2F1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/e2f1-antibody-flow-cytometry-western-blot-elisa/125996>