

Rabbit anti-Human Von Willebrand Factor/VWF Polyclonal Antibody | anti-VWF antibody

Anti-Von Willebrand Factor/VWF Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125999	Rabbit	Human	ELISA, Immunohistochemistry

PRODUCT SYNONYMS

Von Willebrand Factor/VWF; Polyclonal Antibody; Anti-Von Willebrand Factor/VWF Antibody Picoband; anti-VWF antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-VWF antibody	ELISA, IHC (Immunohistochemistry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Von Willebrand Factor/VWF recombinant protein (Position: R1287-Q2770).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of Von Willebrand Factor/VWF using anti-Von Willebrand Factor/VWF antibody (AAA125999). Von Willebrand Factor/VWF was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Von Willebrand Factor/VWF Antibody (AAA125999) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 1. IHC analysis of Von Willebrand Factor/VWF using anti-Von Willebrand Factor/VWF antibody (AAA125999). Von Willebrand Factor/VWF was detected in a paraffin-embedded section of human renal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Von Willebrand Factor/VWF Antibody (AAA125999) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of Von Willebrand Factor/VWF using anti-Von Willebrand Factor/VWF antibody (AAA125999). Von Willebrand Factor/VWF was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Von Willebrand Factor/VWF Antibody (AAA125999) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-VWF antibody	Von Willebrand factor (VWF) is a blood glycoprotein involved in hemostasis. It is mapped to 12p13.31. The VWF gene encodes von Willebrand factor (VWF), a large multimeric glycoprotein that plays a central role in the blood coagulation system, serving both as a major mediator of platelet-vessel wall interaction and platelet adhesion, and as a carrier for coagulation factor VIII. VWF released from endothelial cell Weibel-Palade bodies bound particularly avidly to the extracellular matrix. VWF deficiency or dysfunction (von Willebrand disease) leads to a bleeding tendency, which is most apparent in tissues having high blood flow shear in narrow vessels.
Product Categories/Family for anti-VWF antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	89191868
NCBI GeneID	7450
NCBI Accession #	NP_000543.2
NCBI GenBank Nucleotide #	NM_000552.3
UniProt Accession #	P04275; Q99806
Molecular Weight	309,265 Da
NCBI Official Full Name	von Willebrand factor preproprotein
NCBI Official Synonym Full Names	von Willebrand factor
NCBI Official Symbol	VWF
NCBI Official Synonym Symbols	VWD; F8VWF
NCBI Protein Information	von Willebrand factor; coagulation factor VIII VWF
UniProt Protein Name	von Willebrand factor
UniProt Gene Name	VWF
UniProt Synonym Gene Names	F8VWF; vWF
UniProt Entry Name	VWF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/von-willebrand-factor-vwf-antibody-immunohistochemistry-elisa/125999>