

Rabbit anti-Human CX3CR1 Polyclonal Antibody | anti-CX3CR1 antibody

Anti-CX3CR1 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126000	Rabbit	Human	Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

CX3CR1; Polyclonal Antibody; Anti-CX3CR1 Antibody Picoband; anti-CX3CR1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CX3CR1 antibody	IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human CX3CR1 recombinant protein (Position: M1-H345).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 4. Flow Cytometry analysis of HEL cells using anti-CX3CR1 antibody (AAA126000). Overlay histogram showing HEL cells stained with AAA126000 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CX3CR1 Antibody (AAA126000, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of CX3CR1 using anti-CX3CR1 antibody (AAA126000). CX3CR1 was detected in a paraffin-embedded section of human lymphoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CX3CR1 Antibody (AAA126000) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of CX3CR1 using anti-CX3CR1 antibody (AAA126000). CX3CR1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CX3CR1 Antibody (AAA126000) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 1. IHC analysis of CX3CR1 using anti-CX3CR1 antibody (AAA126000). CX3CR1 was detected in a paraffin-embedded section of human adenocarcinoma of the right colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CX3CR1 Antibody (AAA126000) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-CX3CR1 antibody	CX3C chemokine receptor 1 (CX3CR1) also known as the fractalkine receptor or G-protein coupled receptor 13 (GPR13) is a protein that in humans is encoded by the CX3CR1 gene. Fractalkine is a transmembrane protein and chemokine involved in the adhesion and migration of leukocytes. The protein encoded by this gene is a receptor for fractalkine. The encoded protein also is a coreceptor for HIV-1, and some variations in this gene lead to increased susceptibility to HIV-1 infection and rapid progression to AIDS. Four transcript variants encoding two different isoforms have been found for this gene.
Product Categories/Family for anti-CX3CR1 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503171
NCBI GeneID	1524
NCBI Accession #	NP_001328.1
NCBI GenBank Nucleotide #	NM_001337.3
UniProt Accession #	P49238 ; A0N0N6 ; B2R5Z4 ; J3KP17
Molecular Weight	40,396 Da
NCBI Official Full Name	CX3C chemokine receptor 1 isoform b
NCBI Official Synonym Full Names	chemokine (C-X3-C motif) receptor 1
NCBI Official Symbol	CX3CR1
NCBI Official Synonym Symbols	V28; CCRL1; GPR13; CMKDR1; GPRV28; CMKBRL1
NCBI Protein Information	CX3C chemokine receptor 1; CMK-BRL1; CMK-BRL-1; C-X3-C CKR-1; fractalkine receptor; G protein-coupled receptor 13; G-protein coupled receptor 13; chemokine (C-X3-C) receptor 1; beta chemokine receptor-like 1; chemokine (C-C) receptor-like 1
UniProt Protein Name	CX3C chemokine receptor 1
UniProt Gene Name	CX3CR1
UniProt Synonym Gene Names	CMKBRL1; GPR13; C-X3-C CKR-1; CX3CR1; CMK-BRL1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cx3cr1-antibody-immunohistochemistry-flow-cytometry-elisa/126000>