

Rabbit anti-Mouse, Rat IL-17A/IL17a Polyclonal Antibody | anti-IL17A antibody

Anti-IL-17A/IL17a Antibody Picoband


CATALOG # AAA126021	HOST Rabbit	REACTIVITY Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry
-------------------------------	-----------------------	---------------------------------	--

PRODUCT SYNONYMS


IL-17A/IL17a; Polyclonal Antibody; Anti-IL-17A/IL17a Antibody Picoband; anti-IL17A antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-IL17A antibody	ELISA, IHC (Immunohistochemistry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived rat IL-17A/IL17a recombinant protein (Position: A18-S150).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of IL-17A/IL17a using anti-IL-17A/IL17a antibody (AAA126021).IL-17A/IL17a was detected in a paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IL-17A/IL17a Antibody (AAA126021) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 1. IHC analysis of IL-17A/IL17a using anti-IL-17A/IL17a antibody (AAA126021).IL-17A/IL17a was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IL-17A/IL17a Antibody (AAA126021) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-IL17A antibody	Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.
Product Categories/Family for anti-IL17A antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	2498483
NCBI GeneID	301289
UniProt Accession #	Q61453
Molecular Weight	16,877 Da
NCBI Official Full Name	Interleukin-17A
NCBI Official Synonym Full Names	interleukin 17A
NCBI Official Symbol	IL17a
NCBI Official Synonym Symbols	IL17; IL-17; CTLA-8; IL-17A
NCBI Protein Information	interleukin-17A; Interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8); cytotoxic T-lymphocyte-associated antigen 8
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	IL17a
UniProt Synonym Gene Names	Ctla8; IL17; IL-17; IL-17A; CTLA-8
UniProt Entry Name	IL17_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/il-17a-il17a-antibody-immunohistochemistry-elisa/126021>