

Rabbit anti-Human PPAR gamma/PPARG Polyclonal Antibody | anti-PPARG antibody

Anti-PPAR gamma/PPARG Antibody Picoband


CATALOG # AAA126027	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Flow Cytometry, Functional Assay, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

PPAR gamma/PPARG; Polyclonal Antibody; Anti-PPAR gamma/PPARG Antibody Picoband; anti-PPARG antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-PPARG antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human PPAR gamma/PPARG recombinant protein (Position: H58-K329).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of A431 cells using anti-PPAR Gamma/PPARG antibody (AAA126027).Overlay histogram showing A431 cells stained with AAA126027 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PPAR Gamma/PPARG Antibody (AAA126027, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of PPAR Gamma/PPARG using anti-PPAR Gamma/PPARG antibody (AAA126027).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human placenta tissue lysates,Lane 2: human MCF-7 whole cell lysates,Lane 3: human SK-OV-3 whole cell lysates,Lane 4: human MDA-MB-453 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PPAR Gamma/PPARG antigen affinity purified polyclonal antibody (#AAA126027) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PPAR Gamma/PPARG at approximately 65 kDa. The expected band size for PPAR Gamma/PPARG is at 65 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PPARG antibody	Peroxisome proliferator- activated receptor gamma (PPAR-gamma or PPARG), also known as the glitazone reverse insulin resistance receptor, or NR1C3 (nuclear receptor subfamily 1, group C, member 3) is a type II nuclear receptor (protein regulating genes) that in humans is encoded by the PPARG gene. This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described.
Product Categories/Family for anti-PPARG antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	116284368
NCBI GeneID	5468
NCBI Accession #	NP_005028.4
NCBI GenBank Nucleotide #	NM_005037.5
UniProt Accession #	P37231; O00684; O00710; O14515; Q0QJH8; Q15178; Q15179; Q15180; Q15832; Q86U60; A8K3G6; B5BUA1
Molecular Weight	57,620 Da
NCBI Official Full Name	peroxisome proliferator-activated receptor gamma isoform 1
NCBI Official Synonym Full Names	peroxisome proliferator-activated receptor gamma
NCBI Official Symbol	PPARG

NCBI Official Synonym Symbols	GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARGgamma
NCBI Protein Information	peroxisome proliferator-activated receptor gamma; PPAR gamma; PPAR-gamma; nuclear receptor subfamily 1 group C member 3; peroxisome proliferator-activated receptor gamma 1; peroxisome proliferator-activated nuclear receptor gamma variant 1
UniProt Protein Name	Peroxisome proliferator-activated receptor gamma
UniProt Gene Name	PPARG
UniProt Synonym Gene Names	NR1C3; PPAR-gamma
UniProt Entry Name	PPARG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ppar-gamma-pparg-antibody-flow-cytometry-western-blot-elisa/126027>