

Rabbit anti-Human Clusterin/CLU Polyclonal Antibody | anti-CLU antibody

Anti-Clusterin/CLU Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126046	Rabbit	Human	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


Clusterin/CLU; Polyclonal Antibody; Anti-Clusterin/CLU Antibody Picoband; anti-CLU antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CLU antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Clusterin/CLU recombinant protein (Position: A108-Q320).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of Clusterin/CLU using anti-Clusterin/CLU antibody (AAA126046).Clusterin/CLU was detected in a paraffin-embedded section of human endometrial cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Clusterin/CLU Antibody (AAA126046) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)
Figure 1. Western blot analysis of Clusterin/CLU using anti-Clusterin/CLU antibody (AAA126046).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human placenta tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Clusterin/CLU antigen affinity purified polyclonal antibody (#AAA126046) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Clusterin/CLU at approximately 70 kDa. The expected band size for Clusterin/CLU is at 52 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CLU antibody	The protein encoded by this gene is a secreted chaperone that can under some stress conditions also be found in the cell cytosol. It has been suggested to be involved in several basic biological events such as cell death, tumor progression, and neurodegenerative disorders. Alternate splicing results in both coding and non-coding variants.
Product Categories/Family for anti-CLU antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	355594753
NCBI GeneID	1191
NCBI Accession #	NP_001822.3
NCBI GenBank Nucleotide #	NM_001831.3
UniProt Accession #	P10909; P11380; P11381; Q2TU75; Q5HYC1; Q7Z5B9; B2R9Q1; B3KSE6; AAB25217
Molecular Weight	52,495 Da
NCBI Official Full Name	clusterin preproprotein
NCBI Official Synonym Full Names	clusterin
NCBI Official Symbol	CLU
NCBI Official Synonym Symbols	CLI; AAG4; APOJ; CLU1; CLU2; KUB1; SGP2; APO-J; SGP-2; SP-40; TRPM2; TRPM-2; NA1/NA2
NCBI Protein Information	clusterin; apolipoprotein J; ku70-binding protein 1; sulfated glycoprotein 2; aging-associated protein 4; complement lysis inhibitor; complement cytolysis inhibitor; complement-associated protein SP-40,40; testosterone-repressed prostate message 2
UniProt Protein Name	Clusterin

UniProt Gene Name	CLU
UniProt Synonym Gene Names	APOJ; CLI; KUB1; Apo-J; CLI; TRPM-2
UniProt Entry Name	CLUS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/clusterin-clu-antibody-immunohistochemistry-western-blot-elisa/126046>