

Rabbit anti-Human, Mouse DAO Polyclonal Antibody | anti-DAO antibody

Anti-DAO Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126071	Rabbit	Human, Mouse	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

DAO; Polyclonal Antibody; Anti-DAO Antibody Picoband; anti-DAO antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-DAO antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human DAO recombinant protein (Position: C18-K338).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of U937 cells using anti-DAO antibody (AAA126071).Overlay histogram showing U937 cells stained with AAA126071 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-DAO Antibody (AAA126071, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of DAO using anti-DAO antibody (AAA126071).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HCCT tissue lysates,Lane 2: mouse kidney tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DAO antigen affinity purified polyclonal antibody (#AAA126071) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for DAO at approximately 39 kDa. The expected band size for DAO is at 39 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-DAO antibody	D-amino acid oxidase (DAAO; also OXDA, DAMOX) is an enzyme with the function on a molecular level to oxidize D-amino acids to the corresponding alpha-keto acids, producing ammonia and hydrogen peroxide. This gene encodes the peroxisomal enzyme D-amino acid oxidase. The enzyme is a flavoprotein which uses flavin adenine dinucleotide (FAD) as its prosthetic group. Its substrates include a wide variety of D-amino acids, but it is inactive on the naturally occurring L-amino acids. Its biological function is not known; it may act as a detoxifying agent which removes D-amino acids that accumulate during aging. In mice, it degrades D-serine, a co-agonist of the NMDA receptor. This gene may play a role in the pathophysiology of schizophrenia.
Product Categories/Family for anti-DAO antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	148539837
NCBI GeneID	1610
NCBI Accession #	NP_001908.3
NCBI GenBank Nucleotide #	NM_001917.4
UniProt Accession #	P14920; Q16758; Q8N6R2; B2R7I5
Molecular Weight	39,474 Da
NCBI Official Full Name	D-amino-acid oxidase
NCBI Official Synonym Full Names	D-amino-acid oxidase
NCBI Official Symbol	DAO
NCBI Official Synonym Symbols	DAAO; OXDA; DAMOX

NCBI Protein Information	D-amino-acid oxidase; D-amino acid oxidase
UniProt Protein Name	D-amino-acid oxidase
UniProt Gene Name	DAO
UniProt Synonym Gene Names	DAMOX; DAAO; DAMOX; DAO
UniProt Entry Name	OXDA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/dao-antibody-western-blot-flow-cytometry-elisa/126071>