

Rabbit anti-Human CD163 Polyclonal Antibody | anti-CD163 antibody

Anti-CD163 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126082	Rabbit	Human	Western Blot, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

CD163; Polyclonal Antibody; Anti-CD163 Antibody Picoband; anti-CD163 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CD163 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human CD163 (SWGTVCDSDWDLDDAQVVCQQL).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of JK cells using anti-CD163 antibody (AAA126082).Overlay histogram showing JK cells stained with AAA126082 (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CD163 Antibody (AAA126082, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of CD163 using anti-CD163 antibody (AAA126082).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HCCT tissue lysates,Lane 2: human HCCP tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD163 antigen affinity purified polyclonal antibody (#AAA126082) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CD163 at approximately 150 kDa. The expected band size for CD163 is at 125 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD163 antibody	CD163 (Cluster of Differentiation 163) is a protein that in humans is encoded by the CD163 gene. The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages, and may thereby protect tissues from free hemoglobin-mediated oxidative damage. This protein may also function as an innate immune sensor for bacteria and inducer of local inflammation. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
Product Categories/Family for anti-CD163 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	344179110
NCBI GeneID	9332
NCBI Accession #	NP_004235.4
NCBI GenBank Nucleotide #	NM_004244.5
UniProt Accession #	Q86VB7; Q07898; Q07899; Q07900; Q07901; Q2VLH7; C9JIG2
Molecular Weight	125,451 Da
NCBI Official Full Name	scavenger receptor cysteine-rich type 1 protein M130 isoform a
NCBI Official Synonym Full Names	CD163 molecule
NCBI Official Symbol	CD163
NCBI Official Synonym Symbols	M130; MM130

NCBI Protein Information	scavenger receptor cysteine-rich type 1 protein M130; hemoglobin scavenger receptor; macrophage-associated antigen
UniProt Protein Name	Scavenger receptor cysteine-rich type 1 protein M130
UniProt Gene Name	CD163
UniProt Synonym Gene Names	M130; sCD163
UniProt Entry Name	C163A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd163-antibody-western-blot-flow-cytometry-elisa/126082>