

Rabbit ATF2 Polyclonal Antibody | anti-ATF2 antibody

Anti-ATF2 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126094	Rabbit	Human, Mouse, Rat	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

ATF2; Polyclonal Antibody; Anti-ATF2 Antibody Picoband; anti-ATF2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ATF2 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ATF2 recombinant protein (Position: M1-H454).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of HeLa cells using anti-ATF2 antibody (AAA126094). Overlay histogram showing HeLa cells stained with AAA126094 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ATF2 Antibody (AAA126094, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of ATF2 using anti-ATF2 antibody (AAA126094). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human HepG2 whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human U-87MG whole cell lysates, Lane 5: rat brain tissue lysates, Lane 6: rat thymus tissue lysates, Lane 7: mouse brain tissue lysates, Lane 8: mouse thymus tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ATF2 antigen affinity purified polyclonal antibody (#AAA126094) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ATF2 at approximately 75 kDa. The expected band size for ATF2 is at 55 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ATF2 antibody	ATF2, also known as Activating transcription factor 2, is a protein that in humans is encoded by the ATF2 gene. It is mapped to 2q31.1. This gene encodes a transcription factor that is a member of the leucine zipper family of DNA-binding proteins. This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. The protein forms a homodimer or heterodimer with c-Jun and stimulates CRE-dependent transcription. The protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro, thus, it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. Additional transcript variants have been identified but their biological validity has not been determined.
Product Categories/Family for anti-ATF2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	368711269
NCBI GeneID	1386
NCBI Accession #	NP_001243019.1
NCBI GenBank Nucleotide #	NM_001256090.1
UniProt Accession #	P15336; Q13000; Q4ZFU9; Q53RY2; Q8TAR1; A1L3Z2; D3DPE9
Molecular Weight	54,537 Da
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-2 isoform 1
NCBI Official Synonym Full Names	activating transcription factor 2
NCBI Official Symbol	ATF2
NCBI Official Synonym Symbols	HB16; CREB2; TREB7; CRE-BP1

NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-2; CREB-2; cAMP-dependent transcription factor ATF-2; cAMP-responsive element-binding protein 2; cAMP response element-binding protein CRE-BP1; cyclic AMP-responsive element-binding protein 2; cAMP responsive
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-2
UniProt Gene Name	ATF2
UniProt Synonym Gene Names	CREB2; CREBP1; cAMP-dependent transcription factor ATF-2; CREB-2
UniProt Entry Name	ATF2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/atf2-antibody-western-blot-flow-cytometry-elisa/126094>