

Rabbit EIF4EBP1 Polyclonal Antibody | anti-EIF4EBP1 antibody

Anti-EIF4EBP1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126098	Rabbit	Human, Mouse, Rat	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

EIF4EBP1; Polyclonal Antibody; Anti-EIF4EBP1 Antibody Picoband; anti-EIF4EBP1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-EIF4EBP1 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human EIF4EBP1 (PMEASQSHLRNSPEDK).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of A431 cells using anti-EIF4EBP1 antibody (AAA126098).Overlay histogram showing A431 cells stained with AAA126098 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-EIF4EBP1 Antibody (AAA126098, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of EIF4EBP1 using anti-EIF4EBP1 antibody (AAA126098).EIF4EBP1 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF4EBP1 Antibody (AAA126098) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of EIF4EBP1 using anti-EIF4EBP1 antibody (AAA126098).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HEL whole cell lysates,Lane 2: human PC-3 whole cell lysates,Lane 3: human HEK293 whole cell lysates,Lane 4: rat stomach tissue lysates,Lane 5: rat pancreas tissue lysates,Lane 6: mouse pancreas tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EIF4EBP1 antigen affinity purified polyclonal antibody (#AAA126098) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for EIF4EBP1 at approximately 17 kDa. The expected band size for EIF4EBP1 is at 17 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-EIF4EBP1 antibody

Eukaryotic translation initiation factor 4E-binding protein 1 (also known as 4E-BP1) is a protein that in humans is encoded by the EIF4EBP1 gene. This gene encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758258
NCBI GeneID	1978
NCBI Accession #	NP_004086.1
NCBI GenBank Nucleotide #	NM_004095.3
UniProt Accession #	Q13541; Q6IBN3; B2R502; D3DSW8
Molecular Weight	12,580 Da
NCBI Official Full Name	eukaryotic translation initiation factor 4E-binding protein 1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 4E binding protein 1
NCBI Official Symbol	EIF4EBP1
NCBI Official Synonym Symbols	BP-1; 4EBP1; 4E-BP1; PHAS-I
NCBI Protein Information	eukaryotic translation initiation factor 4E-binding protein 1; eIF4E-binding protein 1; phosphorylated heat- and acid-stable protein regulated by insulin 1
UniProt Protein Name	Eukaryotic translation initiation factor 4E-binding protein 1
UniProt Gene Name	EIF4EBP1
UniProt Synonym Gene Names	4E-BP1; eIF4E-binding protein 1; PHAS-I
UniProt Entry Name	4EBP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/eif4ebp1-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/126098>