

Rabbit nNOS (neuronal)/NOS1 Polyclonal Antibody | anti-NOS1 antibody

Anti-nNOS (neuronal)/NOS1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126104	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS

nNOS (neuronal)/NOS1; Polyclonal Antibody; Anti-nNOS (neuronal)/NOS1 Antibody Picoband; anti-NOS1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Natural and recombinant Mouse SELE
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-NOS1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Samples	Cell culture supernatants, serum and plasma (heparin, citrate)
Detection Range	156 pg/ml - 10,000 pg/ml
Immunogen	E Coli-derived human nNOS (neuronal)/NOS1 recombinant protein (Position: R19-E1320).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of U87 cells using anti-nNOS (neuronal)/NOS1 antibody (AAA126104).Overlay histogram showing U87 cells stained with AAA126104 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-nNOS (neuronal)/NOS1 Antibody (AAA126104, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of nNOS (neuronal)/NOS1 using anti-nNOS (neuronal)/NOS1 antibody (AAA126104).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: rat brain tissue lysates,Lane 2: mouse brain tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-nNOS (neuronal)/NOS1 antigen affinity purified polyclonal antibody (#AAA126104) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for nNOS (neuronal)/NOS1 at approximately 161 kDa. The expected band size for nNOS (neuronal)/NOS1 is at 161 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-NOS1 antibody	Background: E-selectin, also called endothelial-leukocyte adhesion molecule-1 (ELAM-1), is a cell surface glycoprotein expressed by cytokine-activated endothelium that mediates the adhesion of blood neutrophils. An increased expression of E-selectin has been observed in the arterial endothelium interacting with lymphocytes and macrophages in human atherosclerotic lesions. E-selectin plays a critical role in mediating tissue-specific homing of T cells into skin, and of primitive hematopoietic progenitor cells (HPCs) into bone marrow (BM). A structurally and functionally related group of genes, lymph node homing receptor (LHR), granule membrane protein 140 (GMP-140), and endothelial leukocyte adhesion molecule 1 (ELAM-1) are shown to constitute a gene cluster on mouse and human chromosome 1. The standard product used in this kit is recombinant mouse E-Selectin, excluding intercellular E-Selectin and transmembrane domain. It consists of totally 557 amino acids with the molecular mass of 85.8KDa.
Product Categories/Family for anti-NOS1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	10835173
NCBI GeneID	4842
NCBI Accession #	NP_000611.1
NCBI GenBank Nucleotide #	NM_000620.4
UniProt Accession #	P29475; O75713; E9PH30
Molecular Weight	160,970 Da
NCBI Official Full Name	nitric oxide synthase, brain isoform 1
NCBI Official Synonym Full Names	nitric oxide synthase 1 (neuronal)

NCBI Official Symbol	NOS1
NCBI Official Synonym Symbols	NOS; bNOS; nNOS; IHPS1; N-NOS; NC-NOS
NCBI Protein Information	nitric oxide synthase, brain; NOS type I; neuronal NOS; constitutive NOS; peptidyl-cysteine S-nitrosylase NOS1
UniProt Protein Name	Nitric oxide synthase, brain
UniProt Gene Name	NOS1
UniProt Synonym Gene Names	N-NOS; nNOS
UniProt Entry Name	NOS1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/nnos-neuronal-nos1-antibody-flow-cytometry-western-blot-elisa/126104>