

Rabbit anti-Human CD83 Polyclonal Antibody | anti-CD83 antibody

Anti-CD83 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126172	Rabbit	Human	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

CD83; Polyclonal Antibody; Anti-CD83 Antibody Picoband; anti-CD83 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Natural and recombinant Human CCL14
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CD83 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Assay Type	Sandwich
Samples	Cell culture supernatants, cell lysates, serum and plasma (heparin, EDTA)
Detection Range	15.6 pg/ml - 1,000 pg/ml
Sensitivity	<10 pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay): Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision across assays): Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of RT4 cells using anti-ADAM22 antibody (AAA126172).Overlay histogram showing RT4 cells stained with AAA126172 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ADAM22 Antibody (AAA126172, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of ADAM22 using anti-ADAM22 antibody (AAA126172).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Daudi whole cell lysates,Lane 2: human Jurkat whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ADAM22 antigen affinity purified polyclonal antibody (#AAA126172) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ADAM22 at approximately 45 kDa. The expected band size for ADAM22 is at 23 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD83 antibody	Background: Chemokine(C-C motif) ligand 14(CCL14) is a small cytokine belonging to the CC chemokine family. It is also commonly known as HCC-1. It is produced as a protein precursor that is processed to generate a mature active protein containing 74 amino acids that and is 46% identical in amino acid composition to CCL3 and CCL4. This chemokine is expressed in various tissues including spleen, bone marrow, liver, muscle, and gut. CCL14 activates monocytes, but does not induce their chemotaxis. In addition, HCC-1 enhanced the proliferation of CD34+ myeloid progenitor cells. It was as effective as MIP-1 alpha, but about 100-fold less potent. Human CCL14 is located on chromosome 17 within a cluster of other chemokines belonging to the CC family. Principle of the Assay: The Quick Picokine™ Human CCL14 Pre-Coated ELISA (Enzyme-Linked Immunosorbent Assay) kit is a solid phase immunoassay specially designed to measure Human CCL14 in cell culture supernatants, cell lysates, serum and plasma (heparin, EDTA). It uses our proprietary Quick ELISA technology. Quick ELISA technology employs capture antibodies conjugated to an affinity tag that is recognized by the polyclonal antibody used to coat our Quick ELISA plates. This approach to sandwich ELISA allows the formation of the antibody-analyte sandwich complex in a single step, significantly reducing assay time. The kit contains recombinant Human CCL14 with immunogen: Expression system for standard: E.coli; Immunogen sequence: T20-N93. To measure Human CCL14, add standards and samples to the wells, then add antibody cocktail. Wash the wells with PBS or TBS buffer, and add TMB. TMB is an HRP substrate and will be catalyzed to produce a blue color product, which changes into yellow after adding the acidic stop solution. The absorbance of the yellow product is linearly proportional to Human CCL14 in the sample. Read the absorbance of the yellow product in each well using a plate reader, and benchmark the sample wells' readings against the standard curve to determine the concentration of Human CCL14 in the sample.
Product Categories/Family for anti-CD83 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	94421024
NCBI GeneID	9308

NCBI Accession #	NP_001035370.1
NCBI GenBank Nucleotide #	NM_001040280.1
UniProt Accession #	Q01151 ; Q5THX9
Molecular Weight	23,042 Da
NCBI Official Full Name	CD83 antigen isoform b
NCBI Official Synonym Full Names	CD83 molecule
NCBI Official Symbol	CD83
NCBI Official Synonym Symbols	BL11; HB15
NCBI Protein Information	CD83 antigen; hCD83; B-cell activation protein; cell surface protein HB15; cell-surface glycoprotein; CD83 antigen (activated B lymphocytes, immunoglobulin superfamily)
UniProt Protein Name	CD83 antigen
UniProt Gene Name	CD83
UniProt Synonym Gene Names	hCD83
UniProt Entry Name	CD83_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd83-antibody-western-blot-flow-cytometry-elisa/126172>