

Rabbit anti-Human KRT7 Polyclonal Antibody | anti-KRT7 antibody

Anti-KRT7 Antibody Picoband

CATALOG # AAA126225	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

KRT7; Polyclonal Antibody; Anti-KRT7 Antibody Picoband; anti-KRT7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-KRT7 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human KRT7 (QE QKS AKSSRLPDIFE AQIAGLR).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




IHC (Immunohistochemistry)

Figure 4. IHC analysis of KRT7 using anti-KRT7 antibody (AAA126225). KRT7 was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-KRT7 Antibody (AAA126225) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of KRT7 using anti-KRT7 antibody (AAA126225). KRT7 was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-KRT7 Antibody (AAA126225) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of KRT7 using anti-KRT7 antibody (AAA126225). KRT7 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-KRT7 Antibody (AAA126225) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of KRT7 using anti-KRT7 antibody (AAA126225). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HeLa whole cell lysates, Lane 2: human RT4 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-KRT7 antigen affinity purified polyclonal antibody (#AAA126225) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for KRT7 at approximately 51 kDa. The expected band size for KRT7 is at 51 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT7 antibody	Keratin, type II cytoskeletal 7 also known as cytokeratin-7 (CK-7) or keratin-7 (K7) or sarcolectin (SCL) is a protein that in humans is encoded by the KRT7 gene. The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels. The genes encoding the type II cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing may result in several transcript variants; however, not all variants have been fully described.
Product Categories/Family for anti-KRT7 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	67782365
NCBI GeneID	3855
NCBI Accession #	NP_005547.3
NCBI GenBank Nucleotide #	NM_005556.3
UniProt Accession #	P08729; Q92676; Q9BUD8; Q9Y3R7
Molecular Weight	51,386 Da
NCBI Official Full Name	keratin, type II cytoskeletal 7
NCBI Official Synonym Full Names	keratin 7
NCBI Official Symbol	KRT7
NCBI Official Synonym Symbols	K7; CK7; SCL; K2C7
NCBI Protein Information	keratin, type II cytoskeletal 7; CK-7; keratin-7; sarcolectin; cytokeratin 7; cytokeratin-7; type-II keratin Kb7; type II mesothelial keratin K7; keratin, 55K type II cytoskeletal; keratin, simple epithelial type I, K7

UniProt Protein Name	Keratin, type II cytoskeletal 7
UniProt Gene Name	KRT7
UniProt Synonym Gene Names	SCL; CK-7; K7
UniProt Entry Name	K2C7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/krt7-antibody-western-blot-immunohistochemistry-elisa/126225>