

Rabbit anti-Human, Rat ATG4B Polyclonal Antibody | anti-ATG4B antibody

Anti-ATG4B Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126267	Rabbit	Human, Rat	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

ATG4B; Polyclonal Antibody; Anti-ATG4B Antibody Picoband; anti-ATG4B antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ATG4B antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ATG4B recombinant protein (Position: M1-L392).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of Jurkat cells using anti-ATG4B antibody (AAA126267). Overlay histogram showing Jurkat cells stained with AAA126267 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ATG4B Antibody (AAA126267, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of ATG4B using anti-ATG4B antibody (AAA126267). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human 293T whole cell lysates, Lane 3: human HepG2 whole cell lysates, Lane 4: human Jurkat whole cell lysates, Lane 5: human RT4 whole cell lysates, Lane 6: human MCF-7 whole cell lysates, Lane 7: human MOLT-4 whole cell lysates, Lane 8: human HEL whole cell lysates, Lane 9: rat brain tissue lysates, Lane 10: rat liver tissue lysates, Lane 11: rat RH35 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ATG4B antigen affinity purified polyclonal antibody (#AAA126267) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ATG4B at approximately 48 kDa. The expected band size for ATG4B is at 48 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ATG4B antibody	Cysteine protease ATG4B is an enzyme that in humans is encoded by the ATG4B gene. Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
Product Categories/Family for anti-ATG4B antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	47132611
NCBI GeneID	23192
NCBI Accession #	NP_037457.3
NCBI GenBank Nucleotide #	NM_013325.4
UniProt Accession #	Q9Y4P1; Q53NU4; Q6ZUV8; Q8WYM9; Q96K07; Q96K96; Q96SZ1; Q9Y2F2; B7WNK2
Molecular Weight	44,294 Da
NCBI Official Full Name	cysteine protease ATG4B isoform a
NCBI Official Synonym Full Names	autophagy related 4B, cysteine peptidase
NCBI Official Symbol	ATG4B
NCBI Official Synonym Symbols	APG4B; AUTL1
NCBI Protein Information	cysteine protease ATG4B; hAPG4B; autophagin-1; APG4 autophagy 4 homolog B; AUT-like 1 cysteine endopeptidase; ATG4 autophagy related 4 homolog B; autophagy-related protein 4 homolog B; autophagy-related cysteine endopeptidase 1

UniProt Protein Name	Cysteine protease ATG4B
UniProt Gene Name	ATG4B
UniProt Synonym Gene Names	APG4B; AUTL1; KIAA0943; hAPG4B
UniProt Entry Name	ATG4B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/atg4b-antibody-western-blot-flow-cytometry-elisa/126267>