

Rabbit LRIT3 Polyclonal Antibody | anti-LRIT3 antibody

Anti-LRIT3 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127672	Rabbit	Human, Mouse, Rat	ELISA, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

LRIT3; Polyclonal Antibody; Anti-LRIT3 Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-LRIT3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-LRIT3 antibody	ELISA, IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived human LRIT3 recombinant protein (Position: E50-R573).
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



IF (Immunofluorescence)

Figure 2. IF analysis of LTRIT3 using anti-LTRIT3 antibody (AAA127672) and anti-Beta Tubulin antibody (M01857-3).LTRIT3 was detected in immunocytochemical section of U2OS cell. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-LTRIT3 Antibody (AAA127672) and mouse anti-Beta Tubulin antibody (M01857-3) overnight at 4 degree C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) and DyLight488 Conjugated Goat Anti-Mouse IgG (BA1126) were used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37 degree C. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of LTRIT3 using anti-LTRIT3 antibody (AAA127672).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human 293T whole cell lysates,Lane 2: human Jurkat whole cell lysates,Lane 3: rat kidney tissue lysates,Lane 4: mouse kidney tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-LTRIT3 antigen affinity purified polyclonal antibody (#AAA127672) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for LTRIT3 at approximately 75 kDa. The expected band size for LTRIT3 is at 75 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-LTRIT3 antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-LTRIT3 antibody	Primary Antibodies; Signal Transduction; Signaling Pathway; Small G Proteins
References	1. Hartz, P. A. Personal Communication. Baltimore, Md. 1/4/2013.2. Kim, S.-D., Liu, J. L., Roscioli, T., Buckley, M. F., Yagnik, G., Boyadjiev, S. A., Kim, J. Leucine-rich repeat, immunoglobulin-like and transmembrane domain 3 (LRIT3) is a modulator of FGFR1. FEBS Lett. 586: 1516-1521, 2012. 3. Neuille, M., El Shamieh, S., Orhan, E., Michiels, C., Antonio, A., Lancelot, M.-E., Condroyer, C., Bujakowska, K., Poch, O., Sahel, J.-A., Audo, I., Zeitz, C. Lrit3 deficient mouse (nob6): a novel model of complete congenital stationary night blindness (cCSNB). PLoS One 9: e90342, 2014. Note: Electronic Article.

NCBI AND UNIPROT INFORMATION

NCBI GI #	385298685
NCBI GeneID	345193
NCBI Accession #	NP_940908.3
NCBI GenBank Nucleotide #	NM_198506.4
UniProt Accession #	Q3SXY7; Q6ZTG1; C9J1C2
Molecular Weight	59,051 Da
NCBI Official Full Name	leucine-rich repeat, immunoglobulin-like domain and transmembrane domain-containing protein 3
NCBI Official Synonym Full Names	leucine-rich repeat, immunoglobulin-like and transmembrane domains 3

NCBI Official Symbol	LRIT3
NCBI Official Synonym Symbols	CSNB1F; FIGLER4
NCBI Protein Information	leucine-rich repeat, immunoglobulin-like domain and transmembrane domain-containing protein 3; fibronectin type III, immunoglobulin and leucine rich repeat domains 4
UniProt Protein Name	Leucine-rich repeat, immunoglobulin-like domain and transmembrane domain-containing protein 3
UniProt Gene Name	LRIT3
UniProt Entry Name	LRIT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/lrit3-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/127672>