

Rabbit anti-Human PCDHA6 Polyclonal Antibody | anti-PCDHA6 antibody

Anti-PCDHA6 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127808	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS

PCDHA6; Polyclonal Antibody; Anti-PCDHA6 Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-PCDHA6 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-PCDHA6 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Immunogen	E coli-derived human PCDHA6 recombinant protein (Position: D125-Q801).
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of PC-3 cells using anti-PCDHA6 antibody (AAA127808).Overlay histogram showing PC-3 cells stained with AAA127808 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-PCDHA6 Antibody (AAA127808, 1ug/1x106 cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of MCF-7 cells using anti-PCDHA6 antibody (AAA127808).Overlay histogram showing MCF-7 cells stained with AAA127808 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-PCDHA6 Antibody (AAA127808, 1ug/1x106 cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of PCDHA6 using anti-PCDHA6 antibody (AAA127808).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human MCF-7 whole cell lysates,Lane 2: human U2OS whole cell lysates,Lane 3: human PC-3 whole cell lysates,Lane 4: human HepG2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PCDHA6 antigen affinity purified polyclonal antibody (#AAA127808) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PCDHA6 at approximately 150 kDa. The expected band size for PCDHA6 is at 103,120 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PCDHA6 antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-PCDHA6 antibody	Primary Antibodies; Signal Transduction; Signaling Pathway; Small G Proteins

References	1. Ribich, S., Tasic, B., Maniatis, T. Identification of long-range regulatory elements in the protocadherin-alpha gene cluster. Proc. Nat. Acad. Sci. 103: 19719-19724, 2006. 2. Wu, Q., Maniatis, T. A striking organization of a large family of human neural cadherin-like cell adhesion genes. Cell 97: 779-790, 1999. 3. Wu, Q., Zhang, T., Cheng, J.-F., Kim, Y., Grimwood, J., Schmutz, J., Dickson, M., Noonan, J. P., Zhang, M. Q., Myers, R. M., Maniatis, T. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. Genome Res. 11: 389-404, 2001.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	9256592
NCBI GeneID	56142
NCBI Accession #	NP_061732
NCBI GenBank Nucleotide #	NM_018909
UniProt Accession #	Q9UN73; O75283; Q9NRT8
Molecular Weight	86,845 Da
NCBI Official Full Name	protocadherin alpha-6 isoform 1
NCBI Official Synonym Full Names	protocadherin alpha 6
NCBI Official Symbol	PCDHA6
NCBI Official Synonym Symbols	CNR2; CNRN2; CNRS2; CRNR2; PCDH-ALPHA6
NCBI Protein Information	protocadherin alpha-6; KIAA0345-like 8; PCDH-alpha-6
UniProt Protein Name	Protocadherin alpha-6
UniProt Gene Name	PCDHA6
UniProt Synonym Gene Names	CNRS2; PCDH-alpha-6
UniProt Entry Name	PCDA6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/pcdha6-antibody-flow-cytometry-western-blot-elisa/127808>