

Mouse CD20 Monoclonal Antibody | anti-CD20 antibody

Anti-Hu CD20 APC-Cyanine7

CATALOG # AAA128316	HOST Mouse	REACTIVITY Human, Non-human primates	APPLICATIONS Flow Cytometry, Functional Assay
-------------------------------	----------------------	--	---

PRODUCT SYNONYMS

CD20; Monoclonal Antibody; Anti-Hu CD20 APC-Cyanine7; MS4A1; B1; S7; MS4A; Bp35; CVID5; LEU-16; anti-CD20 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Non-human primates
Clonality	Monoclonal
Isotype	IgG2b kappa
Clone Number	2H7
Specificity	The mouse monoclonal antibody 2H7 recognizes an extracellular epitope on CD20 (B1, Bp35), a 33-37 kDa non-glycosylated membrane receptor with four transmembrane domains, expressed on pre-B lymphocytes, resting and activated B cells (not plasma cells), follicular dendritic cells, and at low levels on peripheral blood T lymphocytes.
Purity/Purification	The purified antibody is conjugated with tandem dye APC-Cyanine7 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Form/Format	Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Sequence Length	297
Applicable Applications for anti-CD20 antibody	FCM/FACS (Flow Cytometry)
Immunogen	Human tonsillar B cells
Conjugation	APC-Cyanine7
Preparation and Storage	Store in the dark at 2-8 degree C. Do not freeze. Avoid prolonged exposure to light.

ADDITIONAL INFORMATION

Related Product Information for anti-CD20 antibody	CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on a subpopulation of T lymphocytes and it can be expressed on follicular dendritic cells). Its expression on B cells is synchronous with the expression of surface IgM. CD20 regulates transmembrane calcium conductance (probably functioning as a component of store-operated calcium channel), cell cycle progression and B-cell proliferation. It is associated with lipid rafts, but the intensity of this association depends on extracellular triggering, employing CD20 conformational change and/or BCR (B cell antigen receptor) aggregation. After the receptor ligation, BCR and CD20 colocalize and then rapidly dissociate before BCR endocytosis, whereas CD20 remains at the cell surface. CD20 serves as a useful target for antibody-mediated therapeutic depletion of B cells, as it is expressed at high levels on most B-cell malignancies, but does not become internalized or shed from the plasma membrane following mAb treatment.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	115968
-----------	--------

NCBI GeneID	931
UniProt Accession #	P11836
NCBI Official Full Name	B-lymphocyte antigen CD20
NCBI Official Synonym Full Names	membrane spanning 4-domains A1
NCBI Official Symbol	MS4A1
NCBI Official Synonym Symbols	B1; S7; Bp35; CD20; CVID5; MS4A2; LEU-16
NCBI Protein Information	B-lymphocyte antigen CD20
UniProt Protein Name	B-lymphocyte antigen CD20
UniProt Gene Name	MS4A1
UniProt Synonym Gene Names	CD20
UniProt Entry Name	CD20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/cd20-antibody-clone-2h7-flow-cytometry-elisa/128316>