

Mouse anti-Human CD3 Monoclonal Antibody | anti-CD3E antibody

Anti-Hu CD3 PerCP-Cyanine5.5

CATALOG # AAA128916	HOST Mouse	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay
-------------------------------	----------------------	----------------------------	---

PRODUCT SYNONYMS

CD3; Monoclonal Antibody; Anti-Hu CD3 PerCP-Cyanine5.5; CD3 antigen; epsilon polypeptide; CD3E; T3E; TCRE; anti-CD3E antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2a
Clone Number	OKT3
Specificity	The mouse monoclonal antibody OKT3 recognizes an extracellular epitope on CD3 antigen of the TCR/CD3 complex on mature human T cells. This antibody, also known as Orthoclone OKT3 or Muromonab-CD3, has been extensively used as a drug for therapy of acute, glucocorticoid resistant rejection of allogenic renal, heart and liver transplants. It has also been investigated for use in treating T-cell acute lymphoblastic leukemia.
Purity/Purification	The purified antibody is conjugated with tandem dye PerCP-Cyanine5.5 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Form/Format	Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Sequence Length	207
Applicable Applications for anti-CD3E antibody	FCM/FACS (Flow Cytometry)
Immunogen	Commercially sensitive information
Conjugation	PerCP-Cyanine5.5
Preparation and Storage	Store in the dark at 2-8 degree C. Do not freeze. Avoid prolonged exposure to light.

ADDITIONAL INFORMATION

Related Product Information for anti-CD3E antibody	CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	1345708
-----------	---------

NCBI GeneID	916
UniProt Accession #	P07766
NCBI Official Full Name	T-cell surface glycoprotein CD3 epsilon chain
NCBI Official Synonym Full Names	CD3e molecule
NCBI Official Symbol	CD3E
NCBI Official Synonym Symbols	T3E; TCRE; IMD18
NCBI Protein Information	T-cell surface glycoprotein CD3 epsilon chain
UniProt Protein Name	T-cell surface glycoprotein CD3 epsilon chain
UniProt Gene Name	CD3E
UniProt Synonym Gene Names	T3E
UniProt Entry Name	CD3E_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd3-antibody-clone-okt3-flow-cytometry-elisa/128916>