

Mouse anti-Rat CD4 Monoclonal Antibody | anti-CD4 antibody

Anti-Rt CD4 APC

CATALOG # AAA128925	HOST Mouse	REACTIVITY Rat	APPLICATIONS Flow Cytometry, Functional Assay
-------------------------------	----------------------	--------------------------	---

PRODUCT SYNONYMS

CD4; Monoclonal Antibody; Anti-Rt CD4 APC; CD4 molecule; T4/Leu-3; L3T4; anti-CD4 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Rat
Clonality	Monoclonal
Isotype	IgG2a kappa
Clone Number	OX-35
Specificity	The mouse monoclonal antibody OX-35 reacts with an extracellular epitope of rat CD4 transmembrane glycoprotein (55 kDa).
Purity/Purification	The purified antibody is conjugated with allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Form/Format	Phosphate buffered saline (PBS) solution with 15 mM sodium azide
Concentration	0.5 mg/ml (varies by lot)
Sequence Length	457
Applicable Applications for anti-CD4 antibody	FCM/FACS (Flow Cytometry)
Immunogen	MLR generated rat Th cells
Conjugation	APC
Preparation and Storage	Store in the dark at 2-8 degree C. Do not freeze. Avoid prolonged exposure to light.

ADDITIONAL INFORMATION

Related Product Information for anti-CD4 antibody	CD4 (T4) is a single chain transmembrane glycoprotein and belongs to immunoglobulin supergene family. In extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). Transmembrane region forms 25 aa, cytoplasmic tail consists of 38 aa. Domains 1,2 and 4 are stabilized by disulfide bonds. The intracellular domain of CD4 is associated with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. Extracellular ligands: MHC class II molecules (binds to CDR2-like region in CD4 domain 1); HIV envelope protein gp120 (binds to CDR2-like region in CD4 domain 1); IL-16 (binds to CD4 domain 3), Human seminal plasma glycoprotein gp17 (binds to CD4 domain 1), L-selectin. Intracellular ligands: p56LckCD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection (human immunodeficiency virus; CD4 is primary receptor for HIV-1 surface glycoprotein gp120). CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6978631
NCBI GeneID	24932
NCBI Accession #	NP_036837.1
NCBI GenBank Nucleotide #	NP_036837.1
UniProt Accession #	P05540
NCBI Official Full Name	T-cell surface glycoprotein CD4
NCBI Official Synonym Full Names	Cd4 molecule
NCBI Official Symbol	Cd4
NCBI Official Synonym Symbols	p55; W3/25
NCBI Protein Information	T-cell surface glycoprotein CD4
UniProt Protein Name	T-cell surface glycoprotein CD4
UniProt Gene Name	Cd4
UniProt Entry Name	CD4_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd4-antibody-clone-ox-35-flow-cytometry-elisa/128925>