

Mouse anti-Human CD15 Monoclonal Antibody | anti-CD15 antibody

Anti-Hu CD15 Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA129105	Mouse	Human	Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

CD15; Monoclonal Antibody; Anti-Hu CD15 Purified; Lewis x Blood Group antigen; Le(x); SSEA-1; 3-fucosyl-N-acetyllactosamine; anti-CD15 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Clone Number	W6D3
Specificity	Mouse monoclonal antibody W6D3 recognizes an extracellular epitope of CD15, a cell membrane 3-fucosyl-N-acetyllactosamine (3-FAL) strongly expressed on granulocytes, monocytes, macrophages, mast cells; it is also present on Langerhans cells and some myeloid precursors cells. This is a non-IgM anti-CD15 antibody.
Purity/Purification	Purified by protein-A affinity chromatography
Form/Format	Purified. Phosphate buffered saline (PBS) solution with 15mM sodium azide
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-CD15 antibody	FCM/FACS (Flow Cytometry)
Immunogen	WERI-RB-1 retinoblastoma cell line
Preparation and Storage	Store at 2-8 degree C. Do not freeze. Do not use after expiration date stamped on the label.

ADDITIONAL INFORMATION

Related Product Information for anti-CD15 antibody	CD15 (Lewis X, Le(x); stage specific embryonic antigen-1, SSEA-1) is a trisaccharide determinant (3-fucosyl-N-acetyllactosamine) expressed on several glycolipids, glycoproteins and proteoglycans of various cell types, e.g. granulocytes, mast cells, monocytes, macrophages, cells of gastric mucosa, nervous system or various tumour cells. There are several variants of Lewis x, such as sialyl-Lewis x or sulphated Lewis x. Cells with high surface expression of Le(x) antigen exhibit strong self-aggregation, based on calcium-dependent Le(x)-Le(x) interaction. This process is involved for example in embryo compaction or in autoaggregation of teratocarcinoma cells. Sialyl-Le(x) and its isomer sialyl-Le(a) are ligands of selectins. CD15 expression has been extensively used to confirm identification of Hodgkin's disease.
--	---

References

*Mandalasi M, Dorabawila N, Smith DF, Heimbürg-Molinari J, Cummings RD, Nyame AK: Development and characterization of a specific IgG monoclonal antibody toward the Lewis x antigen using splenocytes of *Schistosoma mansoni*-infected mice. *Glycobiology*. 2013 Jul;23(7):877-92., URL: <https://www.ncbi.nlm.nih.gov/pubmed/23542315>,*Sutherland DR, Ortiz F, Quest G, Illingworth A, Benko M, Nayyar R, Marinov I: High-sensitivity 5-, 6-, and 7-color PNH WBC assays for both Canto II and Navios platforms. *Cytometry B Clin Cytom*. 2018 Jul;94(4):637-651., URL: <https://www.ncbi.nlm.nih.gov/pubmed/29381839>

*Benharroch D, Dima E, Levy A, Ohana-Malka O, Ariad S, Prinsloo I, Mejirovsky E, Sacks M, Gopas J: Differential expression of sialyl and non-sialyl-CD15 antigens on Hodgkin-Reed-Sternberg cells: significance in Hodgkin's disease. *Leuk Lymphoma*. 2000 Sep;39(1-2):185-94., URL: <https://www.ncbi.nlm.nih.gov/pubmedDb=pubmed&Cmd=>,*Hakomori S: Le(X) and related structures as adhesion molecules. *Histochem J*. 1992 Nov;24(11):771-6., URL: <https://www.ncbi.nlm.nih.gov/pubmedDb=pubmed&Cmd=>,*Li C, Wong P, Pan T, Xiao F, Yin S, Chang B, Kang SC, Ironside J, Sy MS: Normal cellular prion protein is a ligand of selectins: binding requires Le(X) but is inhibited by sLe(X). *Biochem J*. 2007 Sep 1;406(2):333-41., URL: <https://www.ncbi.nlm.nih.gov/pubmedDb=pubmed&Cmd=>

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/cd15-antibody-clone-w6d3-flow-cytometry-elisa/129105>