

Rabbit anti-Human CD45 Monoclonal Antibody | anti-CD45 antibody

Anti-CD45 antibody (7E1), IgG1 Chimeric mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA129458	Rabbit	Human	Immunohistochemistry


PRODUCT SYNONYMS

CD45; Monoclonal Antibody; Anti-CD45 antibody (7E1); IgG1 Chimeric mAb; B220; CD45R; GP180; L-CA; LCA; LY5; T200; anti-CD45 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit/Human Fc chimeric IgG1
Clone Number	7E1
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5%-8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Applicable Applications for anti-CD45 antibody	IHC (Immunohistochemistry)
Preparation and Storage	Store at -20 degree C to -80 degree C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80 degree C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

IMAGES



FCM/FACS (Flow Cytometry)

Figure 2. Flow cytometry analysis of antigen binding of anti-human CD45 mAb(AAA129458).



IHC (Immunohistochemistry)

Figure 1. A. AAA129458 at 10ug/ml staining CD45 in Hut78 cells by IHC (AAA129458);B. AAA129458 at 10ug/ml staining CD45 in human tonsil tissue by IHC (AAA129458)

ADDITIONAL INFORMATION

Related Product Information for anti-CD45 antibody	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth; differentiation; mitosis; and oncogenic transformation. This PTP contains an extracellular domain; a single transmembrane segment and two tandem intracytoplasmic catalytic domains; and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes; or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases; and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene; which encode distinct isoforms; have been reported.
Product Categories/Family for anti-CD45 antibody	Monoclonal antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	392307007
NCBI GeneID	5788
NCBI Accession #	NP_002829.3
NCBI GenBank Nucleotide #	NM_002838.4
UniProt Accession #	P08575; Q16614; Q9H0Y6; A8K7W6
Molecular Weight	147,254 Da
NCBI Official Full Name	receptor-type tyrosine-protein phosphatase C isoform 1
NCBI Official Synonym Full Names	protein tyrosine phosphatase, receptor type, C
NCBI Official Symbol	PTPRC
NCBI Official Synonym Symbols	LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180
NCBI Protein Information	receptor-type tyrosine-protein phosphatase C; CD45 antigen; T200 glycoprotein; T200 leukocyte common antigen; protein tyrosine phosphatase, receptor type, c polypeptide
UniProt Protein Name	Receptor-type tyrosine-protein phosphatase C
UniProt Gene Name	PTPRC
UniProt Synonym Gene Names	CD45; L-CA
UniProt Entry Name	PTPRC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd45-antibody-clone-7e1-immunohistochemistry-elisa/129458>