

CD19 recombinant protein

Human CD19 (M75V,R76S,F85S) Protein, hFc Tag

CATALOG #
AAA129497

HOST
HEK293


PRODUCT SYNONYMS

CD19; Human CD19 (M75V; R76S; F85S) Protein; hFc Tag; B4; CVID3; Recombinant human CD19 (M75V; F85S) Protein with C-terminal human Fc tag; CD19 recombinant protein

PRODUCT SPECIFICATIONS


Host	HEK293
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Sequence	CD19(M75V, R76S,F85S) (Pro20-Lys 291)hFc(Gl u99-Ala330)
Tag	C-Human Fc
Preparation and Storage	Store at -20 degree C to -80 degree C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80 degree C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

IMAGES



ELISA

Figure 2. ELISA plate pre-coated by 2 ug/mL (100 uL/well) Human CD19 (M75V,R76S,F85S) Protein, hFc Tag (AAA129497) can bind Anti-CD19(FMC63 biosimilar) mAb in a linear range of 3.20–400 ng/mL.



SDS-PAGE

Figure 1. Human CD19(M75V,R76S,F85S) Protein, hFc Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for CD19 recombinant protein	This gene encodes a member of the immunoglobulin gene superfamily. Expression of this cell surface protein is restricted to B cell lymphocytes. This protein is a reliable marker for pre-B cells but its expression diminishes during terminal B cell differentiation in antibody secreting plasma cells. The protein has two N-terminal extracellular Ig-like domains separated by a non-Ig-like domain, a hydrophobic transmembrane domain, and a large C-terminal cytoplasmic domain. This protein forms a complex with several membrane proteins including complement receptor type 2 (CD21) and tetraspanin (CD81) and this complex reduces the threshold for antigen-initiated B cell activation. Activation of this B-cell antigen receptor complex activates the phosphatidylinositol 3-kinase signalling pathway and the subsequent release of intracellular stores of calcium ions. This protein is a target of chimeric antigen receptor (CAR) T-cells used in the treatment of lymphoblastic leukemia. Mutations in this gene are associated with the disease common variable immunodeficiency 3 (CVID3) which results in a failure of B-cell differentiation and impaired secretion of immunoglobulins. CVID3 is characterized by hypogammaglobulinemia, an inability to mount an antibody response to antigen, and recurrent bacterial infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2020]
Product Categories/Family for CD19 recombinant protein	ECD Proteins

NCBI AND UNIPROT INFORMATION

NCBI GI #	296010921
NCBI GeneID	930
NCBI Accession #	NP_001171569.1
NCBI GenBank Nucleotide #	NM_001178098.1
UniProt Accession #	P15391 ; Q96S68 ; Q9BRD6 ; A0N0P9
Molecular Weight	61,128 Da
NCBI Official Full Name	B-lymphocyte antigen CD19 isoform 1
NCBI Official Synonym Full Names	CD19 molecule
NCBI Official Symbol	CD19
NCBI Official Synonym Symbols	B4; CVID3
NCBI Protein Information	B-lymphocyte antigen CD19; differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte surface antigen B4
UniProt Protein Name	B-lymphocyte antigen CD19
UniProt Gene Name	CD19
UniProt Entry Name	CD19_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/cd19-recombinant-protein-hek293/129497>