

Mouse Nucleolin (NCL) ELISA Kit | NCL elisa kit

Mouse Nucleolin (NCL) ELISA Kit

CATALOG #
AAA129623

REACTIVITY
Mouse


PRODUCT SYNONYMS

Nucleolin (NCL); Mouse Nucleolin (NCL) ELISA Kit; ELISA Kit for Nucleolin (NCL); NCL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of nucleolin. No significant cross-reactivity or interference between nucleolin and analogues was observed.
Samples	Mouse tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	The minimum detectable dose of nucleolin is typically less than 0.059ng/mL. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined by adding two standard deviations to the mean optical density value of twenty zero standard replicates and calculating the corresponding concentration.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level nucleolin were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level nucleolin were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Intended Uses	The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of nucleolin in mouse tissue homogenates, cell lysates and other biological fluids.
Preparation and Storage	For unopened kit, all the reagents should be kept according to the label on each vial. The Standard, Detection Reagent A, Detection Reagent B and the 96-well strip plate should be stored at -20 degree C while others should be at 4 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NCL elisa kit	Principle of the assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to nucleolin. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to nucleolin. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain nucleolin, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of nucleolin in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	55956788
NCBI GeneID	4691
NCBI Official Full Name	nucleolin
NCBI Official Synonym Full Names	nucleolin
NCBI Official Symbol	NCL
NCBI Official Synonym Symbols	C23
NCBI Protein Information	nucleolin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/nucleolin-ncl/129623>