

Interleukin 12A (IL12A) Recombinant Protein | IL12A recombinant protein

Recombinant Interleukin 12A (IL12A)


CATALOG # AAA134276	HOST E.coli	REACTIVITY Homo sapiens (Human)	APPLICATIONS Western Blot, SDS-Page
-------------------------------	-----------------------	---	---

PRODUCT SYNONYMS


Interleukin 12A (IL12A); Recombinant Interleukin 12A (IL12A); IL12A recombinant protein

PRODUCT SPECIFICATIONS

Host	E.coli
Reactivity	Homo sapiens (Human)
Purity/Purification	> 90%
Form/Format	PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.
Concentration	Original Concentration: 50ug/mL (varies by lot)
Sequence	RNLPVATPDP GMFPCLLHSQ NLLRAVSNNL QKARQTLEFY PCTSEEIDHE DITKDKTSTV EACLPLELTK NESCLNSRET SFITNGSCLA SRKTSFMMAL CLSSIYEDLK MYQVEFKTMN AKLLMDPKRQ IFLDQNMLAV IDELMQALNF NSETVPQKSS LEEPDKFYTK IKLCILLHAF RIRAVTIDRV MSYLNAS
Applicable Applications for IL12A recombinant protein	WB (Western Blot), SDS-PAGE, Immunogen, Positive Control
Source	Prokaryotic expression
Residues	Arg23~Ser219
Tags	N-terminal His Tag
Subcellular Location	Secreted
Traits	Freeze-dried powder
Predicted isoelectric point	6.0
Phenomenon explanation	The possible reasons that the actual band size differs from the predicted are as follows: 1. Splice variants: Alternative splicing may create different sized proteins from the same gene. 2. Relative charge: The composition of amino acids may affects the charge of the protein. 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc. 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form. 5. Polymerization of the target protein: Dimerization, multimerization etc.
Usage	Reconstitute in ddH2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



Sequence



SDS-PAGE

NCBI AND UNIPROT INFORMATION

NCBI GI #	24430219
NCBI GeneID	3592
NCBI Accession #	NP_000873.2
NCBI GenBank Nucleotide #	NM_000882.3
UniProt Accession #	P29459; Q96QZ1
Molecular Weight	Predicted Molecular Mass: 29.2kDa Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.
NCBI Official Full Name	interleukin-12 subunit alpha
NCBI Official Synonym Full Names	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)
NCBI Official Symbol	IL12A
NCBI Official Synonym Symbols	P35; CLMF; NFSK; NKSF1; IL-12A
NCBI Protein Information	interleukin-12 subunit alpha; CLMF p35; IL35 subunit; IL-12, subunit p35; interleukin 12, p35; interleukin-12 alpha chain; NF cell stimulatory factor chain 1; NK cell stimulatory factor chain 1; cytotoxic lymphocyte maturation factor 1, p35; cytotoxic lymphocyte maturation factor 35 kDa subunit
UniProt Protein Name	Interleukin-12 subunit alpha
UniProt Gene Name	IL12A
UniProt Synonym Gene Names	NKSF1; IL-12A; CLMF p35; NKSF1
UniProt Entry Name	IL12A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/il12a-recombinant-protein-e-coli-wb-western-blot-sds-page-immunogen-positive-control/134276>