

Human Coagulation Factor II (F2) ELISA Kit | F2 elisa kit

Human Coagulation Factor II (F2) ELISA Kit

CATALOG #
AAA136498

REACTIVITY
Human


PRODUCT SYNONYMS

Coagulation Factor II (F2); Human Coagulation Factor II (F2) ELISA Kit; FII; TM; PT; Thrombin; Prothrombin; Pro-Thrombin; Activation peptide fragment 1; Activation peptide fragment 2; F2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of F2. No significant cross-reactivity or interference between F2 and analogues was observed.
Sequence Length	607
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma and other biological fluids
Detection Range	0.41-300ng/mL
Sensitivity	< 0.17ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level F2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level F2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for F2 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to F2. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to F2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain F2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of F2 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of F2 in human serum, plasma and other biological fluids.
Product Categories/Family for F2 elisa kit	Signal transduction; Hematology

NCBI AND UNIPROT INFORMATION

NCBI GI #	62751950
NCBI GeneID	548514
NCBI Accession #	NP_001015797.1
NCBI GenBank Nucleotide #	NM_001015797.1
NCBI Official Full Name	prothrombin
NCBI Official Synonym Full Names	coagulation factor 2 (thrombin)
NCBI Official Symbol	f2
NCBI Protein Information	prothrombin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.