

Human Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit | VCAM1 elisa kit

Human Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit

CATALOG #
AAA14024

REACTIVITY
Human


PRODUCT SYNONYMS

Vascular Cell Adhesion Molecule 1 (VCAM1); Human Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit; CD106 INCAM-100 L1CAM; VCAM1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of VCAM1. No significant cross-reactivity or interference between VCAM1 and analogues was observed.
Sequence Length	739
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Detection Range	0.781-50ng/mL
Sensitivity	< 0.33ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level VCAM1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level VCAM1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VCAM1 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to VCAM1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to VCAM1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain VCAM1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of VCAM1 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of VCAM1 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507875
NCBI GeneID	7412
NCBI Accession #	NP_001069.1
NCBI GenBank Nucleotide #	NM_001078.3
UniProt Accession #	P19320; Q6NUP8; A8K6R7; B4DKS4; E9PDD1
Molecular Weight	74,346 Da
NCBI Official Full Name	vascular cell adhesion protein 1 isoform a
NCBI Official Synonym Full Names	vascular cell adhesion molecule 1
NCBI Official Symbol	VCAM1
NCBI Official Synonym Symbols	CD106; INCAM-100
NCBI Protein Information	vascular cell adhesion protein 1
UniProt Protein Name	Vascular cell adhesion protein 1
UniProt Gene Name	VCAM1
UniProt Synonym Gene Names	L1CAM; V-CAM 1; VCAM-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/vascular-cell-adhesion-molecule-1-vcam1/14024>