

Rat Interleukin 2 (IL2) CLIA Kit | IL2 clia kit

Rat Interleukin 2 (IL2) CLIA Kit

CATALOG #
AAA141037

REACTIVITY
Rat


PRODUCT SYNONYMS

Interleukin 2 (IL2); Rat Interleukin 2 (IL2) CLIA Kit; TCGF; Lymphokine; T-Cell Growth Factor; Aldesleukin; IL2 clia kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Interleukin 2 (IL2). No significant cross-reactivity or interference between Interleukin 2 (IL2) and analogues was observed.
Sequence Length	153
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	1.37-1,000pg/mL
Sensitivity	0.57pg/mL
Application	Chemiluminescent immunoassay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Interleukin 2 (IL2) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Interleukin 2 (IL2) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL2 clia kit	The microplate provided in this kit has been pre-coated with an antibody specific to Interleukin 2 (IL2). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Interleukin 2 (IL2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Interleukin 2 (IL2) level in the sample or standard.;
Product Categories/Family for IL2 clia kit	Cytokine; Infection immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	28178861
NCBI GeneID	3558
NCBI Accession #	NP_000577.2
NCBI GenBank Nucleotide #	NM_000586.3
Molecular Weight	17,628 Da
NCBI Official Full Name	interleukin-2
NCBI Official Synonym Full Names	interleukin 2
NCBI Official Symbol	IL2
NCBI Official Synonym Symbols	IL-2; TCGF; lymphokine
NCBI Protein Information	interleukin-2
UniProt Protein Name	Interleukin-2
UniProt Gene Name	IL2
UniProt Synonym Gene Names	IL-2; TCGF
UniProt Entry Name	IL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.