

Human Osteoprotegerin (OPG) CLIA Kit | OPG clia kit

Human Osteoprotegerin (OPG) CLIA Kit

CATALOG #
AAA141138

REACTIVITY
Human


PRODUCT SYNONYMS

Osteoprotegerin (OPG); Human Osteoprotegerin (OPG) CLIA Kit; TNFRSF11B; TNFRSF11-B; OCIF; OPG; TR1; Tumor Necrosis Factor Receptor Superfamily Member 11b; Osteoclastogenesis Inhibitory Factor; OPG clia kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Osteoprotegerin (OPG). No significant cross-reactivity or interference between Osteoprotegerin (OPG) and analogues was observed.
Sequence Length	401
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	41.2-30,000pg/mL
Sensitivity	14.9pg/mL
Application	Chemiluminescent immunoassay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Osteoprotegerin (OPG) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Osteoprotegerin (OPG) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for OPG clia kit	The microplate provided in this kit has been pre-coated with an antibody specific to Osteoprotegerin (OPG). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Osteoprotegerin (OPG). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Osteoprotegerin (OPG) level in the sample or standard.;
Product Categories/Family for OPG clia kit	Cytokine; Metabolic pathway; Bone metabolism

NCBI AND UNIPROT INFORMATION

NCBI GI #	2072185
NCBI GeneID	4982
Molecular Weight	46,026 Da
NCBI Official Full Name	osteoprotegerin
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 11b
NCBI Official Symbol	TNFRSF11B
NCBI Official Synonym Symbols	OPG; TR1; OCIF; PDB5
NCBI Protein Information	tumor necrosis factor receptor superfamily member 11B
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 11B
UniProt Gene Name	TNFRSF11B
UniProt Synonym Gene Names	OCIF; OPG
UniProt Entry Name	TR11B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-clia-kits/osteoprotegerin-opg/141138>