

Rabbit anti-Mouse, Rat B-Cell Activation Factor Receptor Polyclonal Antibody | anti-BAFFR antibody

B-Cell Activation Factor Receptor (BAFFR) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA144752	Rabbit	Mouse, Rat	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

B-Cell Activation Factor Receptor; Polyclonal Antibody; B-Cell Activation Factor Receptor (BAFFR) Polyclonal Antibody; CD268; TNFRSF13C; TNFRSF13-C; BAFFR; BAFFR; BR3; Tumor Necrosis Factor Receptor Superfamily Member 13C; anti-BAFFR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against BAFFR. It has been selected for its ability to recognize BAFFR in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	500ug/mL (varies by lot)
Applicable Applications for anti-BAFFR antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Rat
Organism Species	Rat
Fragment	BAFFR (Gln21~Lys169?)
Quality Control	Content: The quality control contains recombinant BAFFR (Gln21~Lys169 ^{tt}) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescient (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.


IMAGES



IHC (Immunohistochemisry)
DAB staining on fromalin fixed paraffin- embedded Kidney tissue



WB (Western Blot)
Western Blot: Sample: Recombinant protein.



WB (Western Blot)
Western Blot: Sample: Rat Spleen lysate; Primary Ab: 3ug/ml Rabbit Anti-Rat BAFFR
Antibody Second Ab: 0.2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

UniProt Accession #	D3ZNT3
---------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/b-cell-activation-factor-receptor-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/144752>