

Interleukin 6 (IL6) Active Protein | IL6 active protein

Active Interleukin 6 (IL6)

CATALOG # AAA146609	HOST 293F cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---


PRODUCT SYNONYMS

Interleukin 6 (IL6); Active Interleukin 6 (IL6); MGI2-A; MGI2A; HGF; BSF2; HSF; IFNB2; B-Cell Stimulatory Factor-2; Hybridoma/Plasmacytoma Growth Factor; Hepatocyte Stimulating Factor; Cytotoxic T-Cell Differentiation Factor; IL6 active protein

PRODUCT SPECIFICATIONS

Host	293F cell
Purity/Purification	> 97%
Form/Format	Freeze-dried powder
Sequence Length	241
Applicable Applications for IL6 active protein	Activity Assay, Cell Culture
Fragment	Val30~Met212
Tag	N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Organism Species	Human
Isoelectric Point	6.2
Usage	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Preparation and Storage	Avoid repeated freeze/thaw cycles. Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



Application Data






Figure 2. The inhibitory effect of IL-6 on cell proliferation of MCF-7 cells detected by CCK8.

Application Data




Application Data



Activity

Interleukin 6 (IL-6) is an interleukin that acts as both a pro-inflammatory cytokine and an anti-inflammatory myokine. Current data suggest that direct application of IL-6 on breast cancer cells inhibits proliferation in ER-positive (estrogen- receptor- positive) cells through the Jak/Stat3 pathway. To test the inhibitory effect of IL-6 on proliferation of ER-positive MCF-7 cell line, cells were seeded into triplicate wells of 96-well plates at a density of 5,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of IL-6. After incubated for 96h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10uL of CCK-8 solution was added to each well of the plate, then measure the absorbance at 450nm using a microplate reader after incubating the plate for 1-4 hours at 37°C. The inhibitory effect of IL-6 on cell proliferation of MCF-7 cells observed by inverted microscope and detected by CCK-8 was shown in Figure 1 and Figure 2 respectively (Dose-dependent effect was not detected in this case).

SDS-PAGE



ADDITIONAL INFORMATION

Product Categories/Family for IL6 active protein	Active Protein
---	----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	45382889
NCBI GeneID	395337
NCBI Accession #	NP_989959.1
NCBI GenBank Nucleotide #	NM_204628.1
UniProt Accession #	Q90Y10
Molecular Weight	22.4kDa; 22/24kDa
NCBI Official Full Name	interleukin-6
NCBI Official Synonym Full Names	interleukin 6
NCBI Official Symbol	IL6
NCBI Official Synonym Symbols	IL-6; CHIL-6; interleukin-6

NCBI Protein Information	interleukin-6
UniProt Protein Name	Interleukin-6
UniProt Gene Name	IL6
UniProt Synonym Gene Names	ChIL-6; IL-6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/293f-cell-active-protein/interleukin-6-il6/146609>