

General Glycine (Gly) ELISA Kit | Gly elisa kit

General Glycine (Gly) ELISA Kit

CATALOG #
AAA148433REACTIVITY
General

PRODUCT SYNONYMS

Glycine (Gly); General Glycine (Gly) ELISA Kit; Gly elisa kit

PRODUCT SPECIFICATIONS

Reactivity	General
Specificity	This assay has high sensitivity and excellent specificity for detection of glycine. No significant cross-reactivity or interference between glycine and analogues was observed.
Assay Type	Quantitative Competitive
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	1.23-100ug/mL
Sensitivity	< 0.59ug/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level glycine were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level glycine were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

ADDITIONAL INFORMATION

Related Product Information for Gly elisa kit

Principle of the Assay: This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to glycine has been pre-coated onto a microplate. A competitive inhibition reaction is launched between biotin labeled glycine and unlabeled glycine (Standards or samples) with the pre-coated antibody specific to glycine. After incubation the unbound conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. The amount of bound HRP conjugate is reverse proportional to the concentration of glycine in the sample. After addition of the substrate solution, the intensity of color developed is reverse proportional to the concentration of glycine in the sample.

Intended Uses: The kit is a competitive inhibition enzyme immunoassay technique for the in vitro quantitative measurement of glycine in serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.comOnline datasheet: <https://www.aaabiotech.com/general-elisa-kits/glycine-gly/148433>