

Interleukin 10 Active Protein | IL10 active protein

Active Interleukin 10 (IL10)

CATALOG #
AAA150055



APPLICATIONS
Activity Assay, Cell Culture

PRODUCT SYNONYMS

Interleukin 10; Active Interleukin 10 (IL10); CSIF; IL10A; TGIF; Cytokine Synthesis Inhibitory Factor; IL10 active protein


PRODUCT SPECIFICATIONS

Purity/Purification	>97%
Form/Format	Freeze-dried powder 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Sequence Positions	Ser19~Ser178
Sequence Length	178
Applicable Applications for IL10 active protein	Activity Assay, Cell Culture
Species	Mouse
Tag	N-terminal His Tag
Isoelectric Point	8.2
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (PH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Preparation and Storage	Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. Avoid repeated freeze/thaw cycles. Stability: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Gene Sequencing
Figure. Gene Sequencing (Extract

Bioactivity
Figure. The binding activity of IL-10 with IL10Ra. Interleukin 10 (IL-10), also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. IL-10 is a cytokine with multiple, pleiotropic, effects in immunoregulation and inflammation. It downregulates the expression of Th1 cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. IL-10 can block NF- κ B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Besides, Interleukin 10 Receptor Alpha (IL10Ra) has been identified as an interactor of IL-10, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse IL-10 and recombinant mouse IL10Ra. Briefly, IL-10 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100L were then transferred to IL10Ra-coated microtiter wells and incubated for 2h at 37. Wells were washed with PBST and incubated for 1h with anti-IL-10 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37. Finally, add 50uL stop solution to the wells and read at 450nm immediately. The binding activity of IL-10 and IL10Ra was shown in Figure 1, and this effect was in a dose dependent manner.



WB (Western Blot)
Figure. Western Blot



SDS-PAGE
Figure. SDS-PAGE

ADDITIONAL INFORMATION

Product Categories/Family for IL10 active protein	Cytokine; Infection immunity
---	------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	6754318
NCBI GeneID	16153
NCBI Accession #	NP_034678.1
NCBI GenBank Nucleotide #	NP_034678.1
UniProt Accession #	P18893

Molecular Weight	Predicted Molecular Mass: 20.4kDa Accurate Molecular Mass: 22kDa
NCBI Official Full Name	interleukin-10
NCBI Official Synonym Full Names	interleukin 10
NCBI Official Symbol	Il10
NCBI Official Synonym Symbols	CSIF; IL-10
NCBI Protein Information	interleukin-10
UniProt Protein Name	Interleukin-10
UniProt Gene Name	Il10
UniProt Synonym Gene Names	IL-10; IL-10; CSIF
UniProt Entry Name	IL10_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/il10-active-protein/interleukin-10/150055>