

Angiotensin I Converting Enzyme 2 (ACE2) Active Protein | ACE2 active protein

Active Angiotensin I Converting Enzyme 2 (ACE2)

CATALOG # AAA150976	HOST 293F Cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---

PRODUCT SYNONYMS

Angiotensin I Converting Enzyme 2 (ACE2); Active Angiotensin I Converting Enzyme 2 (ACE2); ACE2/Angiotensin I Converting Enzyme 2; ACEH; ACEII; ACE-II; Peptidyl-Dipeptidase A; ACE-related carboxypeptidase; Angiotensin-converting enzyme homolog; Metalloprotease MPROT15; ACE2 active protein

PRODUCT SPECIFICATIONS

Host	293F Cell
Purity/Purification	Greater than 95% by SDS-PAGE
Form/Format	Freeze-dried powder 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Sequence	Gln18~Ser7 40
Applicable Applications for ACE2 active protein	Activity Assay, Cell Culture
Endotoxin	<1.0EU per 1ug (determined by the LAL method)
Tag	N-terminal His Tag
Biological Activity	Active
Expression System	Prokaryotic expression
Organism Species	Human
Isoelectric Point	5.1
Subcellular Location	Secreted
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0mg/mL. Do not vortex.
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months Stable for 12 months if correctly stored. The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

```
QST IEEQAKTFLD KFNHEAEDLF YQSSLASWNY
NTNITEENVQ NMNAGDKWS AFLKEQSTLA QMYPLQEIQN LTVKLQLQAL
QQNGSSVLSE DKSRLNTIL NTMSTIYSTG KVCNPDNPQE CLLLEPGLNE
IMANSLDYNE RLWAWESWRS EVGKQLRPLY EEYVVLKNEM ARANHYEDYG
DYWRGDYEVN GVDGYDYSRG QLIEDVEHTF EEIKPLYEHL HAYVRAKLMN
AYPSYISPIG CLPAHLLGDM WGRFWNLYS LTVPFQKPN IDVTAMVDQ
AWDAQRIFKE AEKFFVSVGL PNMTQGFWEH SMLTDPGNVQ KAVCHPTAHD
LGKGDFRILM CTKVTMDDFL TAHHEMGHIQ YDMAYAAQPF LLRNGANEGF
HEAVGEIMSL SAATPKHLKS IGLLSPDFQE DNETEINFLL KQALTIIVGTL
PFTYMLEKWR WMVFKEIPK DQWMMKWMEM KREIVGVVEP VPHDETYCDP
ASLFHVSNDY SFIRYYTRL YQFQFQALC QAAKHEGPLH KCDISNSTEA
GQKLFNMLRL GKSEPTLAL ENNVGAKNMN VRPLLNYFEP LFTWLKDQNK
NSFVGWSTDW SPYADQSIKV RISLKSALGD KAYEWNNDNM YLFRSSVAYA
MRQYFLKVKV QMILFGEDV RVANLKPRIS FNFFVTAPKN VSDIIPRTEV
EKAIRMSRSR INDAFRLNDN SLEFLGIQPT LGPPNQPPVS
```

Sequence






Figure 1. The binding activity of ACE2 with RBD

Bioactivity


Angiotensin-converting enzyme 2 (ACE2), as a transmembrane protein, serves as the main entry point into cells for some coronaviruses, including HCoV-NL63, SARS-CoV , and SARS-CoV-2 . More specifically, the binding of the spike S1 protein of SARS-CoV and SARS-CoV-2 to the enzymatic domain of ACE2 on the surface of cells results in endocytosis and translocation of both the virus and the enzyme into endosomes located within cells. Besides, recent studies show that spike (S) proteins of 2019-nCoV and SARS-CoV may use the same host cell receptor called angiotensin-converting enzyme 2 (ACE2) for entering into host cells, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ACE2 and recombinant Spike glycoprotein RBD. Briefly, biotin-linked ACE2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100µl were then transferred to RBD-coated microtiter wells and incubated for 2h at 37?. Wells were washed with PBST 3 times and incubation with HRP conjugate for 30min, then wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37?. Finally, add 50ul stop solution to the wells and read at 450nm immediately. The binding activity of ACE2 and RBD was shown in Figure 1, and this effect was in a dose dependent manner.



WB (Western Blot)
WB Image



SDS-PAGE
SDS-PAGE



Sequence
Gene Sequencing (extract)

ADDITIONAL INFORMATION

Related Product Information for ACE2 active protein	Angiotensin-converting-enzyme-2(ACE2),as a transmembrane protein, serves as the main entry point into cells for some coronaviruses,including HCoV-NL63, SARS-CoV , and SARS-CoV-2 .More specifically, the binding of the spike S1 protein of SARS-CoV and SARS-CoV-2 to theenzymatic domain of ACE2 on the surface of cells results in endocytosis and translocation of both the virusand the enzyme into endosomes located within cells. Besides, recent studies show that spike (S) proteins of 2019-nCoV andSARS-CoV may use the same host cell receptor called angiotensin-convertingenzyme 2 (ACE2) for entering into host cells, thus a binding ELISA assay wasconducted to detect the interaction of recombinant human ACE2 andrecombinant Spike glycoprotein RBD. Briefly, biotin-linked ACE2 were dilutedserially in PBS, with 0.01% BSA (pH7.4). Duplicate samples of 100ul were thentransferred to RBD-coated microtiter wells and incubated for 2h at 37 degree C. Wellswere washed with PBST 3 times and incubation with HRP conjugage for 30min, then wells were aspirated and washed 5 times. With the addition of substratesolution, wells were incubated 15-25 minutes at 37 degree C. Finally, add 50ul stopsolution to the wells and read at 450nm immediately. The binding activity of ACE2 and RBD was shown in Figure 1, and this effect was in a dose dependent manner.
Product Categories/Family for ACE2 active protein	Enzyme & Kinase; Metabolic pathway; Endocrinology; Cardiovascular biology; Neuro science; Hepatology; Active, eukaryotic recombinant

NCBI AND UNIPROT INFORMATION

NCBI GI #	11225609
NCBI GeneID	59272
NCBI Accession #	NP_068576.1
NCBI GenBank Nucleotide #	NM_021804.2
UniProt Accession #	Q9BYF1; Q6UWP0; Q86WT0; Q9NRA7; Q9UFZ6; C7ECU1
Molecular Weight	92,463 Da
NCBI Official Full Name	angiotensin-converting enzyme 2
NCBI Official Synonym Full Names	angiotensin I converting enzyme 2
NCBI Official Symbol	ACE2
NCBI Official Synonym Symbols	ACEH
NCBI Protein Information	angiotensin-converting enzyme 2; peptidyl-dipeptidase A; metalloprotease MPROT15; ACE-related carboxypeptidase; angiotensin-converting enzyme homolog; angiotensin I converting enzyme (peptidyl-dipeptidase A) 2
UniProt Protein Name	Angiotensin-converting enzyme 2
UniProt Gene Name	ACE2
UniProt Synonym Gene Names	ACEH
UniProt Entry Name	ACE2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/293f-active-protein/angiotensin-i-converting-enzyme-2-ace2/150976>