

E-cadherin active protein

Active E-cadherin


CATALOG # AAA161657	HOST 293F cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---

PRODUCT SYNONYMS

E-cadherin; Active E-cadherin; E-cadherin active protein (eukaryotic; 293F cell); Human; CDHE; CD324; CDH1; Arc1; ECAD; CDH-E; E-CAD; LCAM; UVO; Cadherin 1 Type 1; Uvomorulin; E-Cadherin; Calcium-Dependent Adhesion Protein; Epithelial; Liver Cell Adhesion Molecule4; E-cadherin active protein


PRODUCT SPECIFICATIONS

Host	293F cell
Purity/Purification	Greater than 90% by SDS-PAGE
Form/Format	Freeze-dried powder; PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Sequence	Asp155~Ile 707
Applicable Applications for E-cadherin active protein	Activity Assay, Cell Culture
Species	Human
Expression System	Eukaryotic expression
Tag	N-terminal His Tag
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Isoelectric Point	4
Research Area	CD & Adhesion molecule, Tumor immunity
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months



SDS-PAGE

Figure. SDS-PAGE



Bioactivity
E-Cadherin, also known as Uvomorulin in the mouse and rat, is a 120 kDa member of the Cadherin family of cell surface glycoproteins that mediate cell adhesion. Human E-Cadherin shares 81% amino acid sequence identity with the rat and mouse proteins. It is a single-pass transmembrane protein that mediates calcium-dependent epithelial cell adhesion. As E-cadherin has the function of cell adhesion, we measure the activity of recombinant human E-cadherin by the ability of the immobilized protein to support the adhesion of MCF7 human breast cancer cells. When 5 x 10⁴ cells/well are added to different concentrations of recombinant human E-cadherin coated plates, cells will adhere after 2 hour incubation at 37 degree C. The adhesion of MCF7 after 2 hour incubation at 37 degree C observed by inverted microscope was shown in Figure 1. Cell adherent was in a dose dependent manner, the result was shown in Figure 2, the EC50 was 0.23 ug/ml.

ADDITIONAL INFORMATION

Product Categories/Family for E-cadherin active protein	Active proteins
---	-----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757960
NCBI GeneID	999
NCBI Accession #	NP_004351.1
NCBI GenBank Nucleotide #	NM_004360.3
UniProt Accession #	P12830; Q13799; Q14216; Q15855; Q16194; Q4PJ14; A8K1U7
Molecular Weight	97,456 Da
NCBI Official Full Name	cadherin-1 preproprotein
NCBI Official Synonym Full Names	cadherin 1, type 1, E-cadherin (epithelial)
NCBI Official Symbol	CDH1
NCBI Official Synonym Symbols	UVO; CDHE; ECAD; LCAM; Arc-1; CD324
NCBI Protein Information	cadherin-1; CAM 120/80; E-Cadherin; uvomorulin; cell-CAM 120/80; epithelial cadherin; cadherin 1, E-cadherin (epithelial); calcium-dependent adhesion protein, epithelial
UniProt Protein Name	Cadherin-1
UniProt Gene Name	CDH1
UniProt Synonym Gene Names	CDHE; UVO; E-cadherin
UniProt Entry Name	CADH1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/293f-active-protein/e-cadherin/161657>