

Porcine CFTR ELISA Kit | CFTR elisa kit

Porcine CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) ELISA Kit

CATALOG #
AAA165426

REACTIVITY
Porcine


PRODUCT SYNONYMS

CFTR; Porcine CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) ELISA Kit; CFTR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of Porcine CFTR. No significant cross-reactivity or interference between Porcine CFTR and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross- reactivity detection between Porcine CFTR and all the analogues, therefore, cross reaction may still exist.
Sequence Length	13
Samples	Serum, Plasma, Biological Fluids
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	Min: 0.094ng/mL; Max: 10ng/mL
Preparation and Storage	Store at 4 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CFTR elisa kit	Description: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CFTR in Porcine serum, plasma and other biological fluids Principle of the Assay: This ELISA kit uses Sandwich-ELISA as the method. The micro ELISA plate provided in this kit has been pre-coated with an antibody specific to Porcine CFTR. Standards or samples are added to the appropriate micro ELISA plate wells and bound by the specific antibody. Then a biotinylated detection antibody specific for Porcine CFTR and Avidin-Horseradish Peroxidase (HRP) conjugate is added to each micro plate well successively and incubated. Free components are washed away. The substrate solution is added to each well. Only those wells that contain Porcine CFTR, biotinylated detection antibody and Avidin-HRP conjugate will appear blue in color. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the color turns yellow. The optical density (OD) is measured spectrophotometrically at a wavelength of 450 nm +/- 2 nm. The OD value is proportional to the concentration of Porcine CFTR. You can calculate the concentration of Porcine CFTR in the samples by comparing the OD of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	180286
NCBI GeneID	1080
UniProt Accession #	P13569; Q20BG8; Q20BH2; Q2I0A1; Q2I102
Molecular Weight	69,230 Da
NCBI Official Full Name	CFTR, partial
NCBI Official Synonym Full Names	cystic fibrosis transmembrane conductance regulator (ATP-binding cassette sub-family C, member 7)
NCBI Official Symbol	CFTR
NCBI Official Synonym Symbols	CF; MRP7; ABC35; ABCC7; CFTR/MRP; TNR-CFTR; dJ760C5.1
NCBI Protein Information	cystic fibrosis transmembrane conductance regulator; cAMP-dependent chloride channel; channel conductance-controlling ATPase
UniProt Protein Name	Cystic fibrosis transmembrane conductance regulator
UniProt Gene Name	CFTR
UniProt Synonym Gene Names	ABCC7; CFTR
UniProt Entry Name	CFTR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.