

Rabbit ARC Polyclonal Antibody | anti-ARC antibody

ARC Polyclonal Antibody

CATALOG # AAA167962	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

ARC; Polyclonal Antibody; ARC Polyclonal Antibody; Arg3.1; anti-ARC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1.1mg/mL (varies by lot)
Sequence Length	396
Applicable Applications for anti-ARC antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human ARC
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human gastric cancer using ARC Polyclonal Antibody at dilution of 1:25



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human thyroid cancer using ARC Polyclonal Antibody at dilution of 1:25



WB (Western Blot)

Western Blot analysis of Human lymphoma tissue using ARC Polyclonal Antibody at dilution of 1:200

ADDITIONAL INFORMATION

Related Product Information for anti-ARC antibody	Arc (for activity-regulated cytoskeleton-associated protein) is a growth factor and immediate early gene that is enriched in brain. Arc mRNA and protein levels are induced by neuronal activity, which is necessary to stimulate neuroplasticity, indicating a potential role for Arc in activity-dependent changes in dendrite function. Arc expression has been detected in neuronal cell bodies and dendrites in the hippocampus, amygdala, hypothalamus, striatum and cortex. Arc has been shown to localize to the cytoskeleton of neuronal cells and appears to colocalize with F-Actin, although it may associate with an Actin-associated protein rather than directly with F-Actin. It has been shown that cocaine-stimulated neuronal activity results in increased Arc mRNA levels in striatum.
Product Categories/Family for anti-ARC antibody	Neuroscience

NCBI AND UNIPROT INFORMATION

NCBI GI #	8567330
NCBI GeneID	23237
NCBI Accession #	NP_056008.1
NCBI GenBank Nucleotide #	NM_015193.4
Molecular Weight	45,316 Da
NCBI Official Full Name	activity-regulated cytoskeleton-associated protein
NCBI Official Synonym Full Names	activity-regulated cytoskeleton-associated protein
NCBI Official Symbol	ARC
NCBI Official Synonym Symbols	Arg3.1
NCBI Protein Information	activity-regulated cytoskeleton-associated protein; ARC; ARC/ARG3.1; activity-regulated gene 3.1 protein homolog
UniProt Protein Name	Activity-regulated cytoskeleton-associated protein
UniProt Gene Name	ARC
UniProt Synonym Gene Names	Arg3.1
UniProt Entry Name	ARC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/arc-antibody-immunohistochemistry-western-blot-elisa/167962>